

# Elastase-2B siRNA (h): sc-105327

## BACKGROUND

Elastase-2, also known as Elastase-2A, CELA2A (chymotrypsin-like elastase family, member 2A) or ELA2A, is a 269 amino acid secreted protein that belongs to the peptidase S1 family and contains one peptidase S1 domain. Expressed in pancreatic tissue, Elastase-2 interacts with CPA1 and catalyzes the hydrolysis of elastin, specifically cleaving the Leu-[Xaa, Met-[Xaa and Phe-[Xaa residues within elastin. Elastase-2B, like Elastase-2, is a 269 amino acid protein that catalyzes the hydrolysis of elastin. The genes encoding Elastase-2 and Elastase-2B map to human chromosome 1, which spans 260 million base pairs, contains over 3,000 genes and comprises nearly 8% of the human genome. Chromosome 1 houses a large number of disease-associated genes, including those that are involved in familial adenomatous polyposis, Stickler syndrome, Parkinson's disease, Gaucher disease, schizophrenia and Usher syndrome. Aberrations in chromosome 1 are found in a variety of cancers, including head and neck cancer, malignant melanoma and multiple myeloma.

## REFERENCES

1. Fletcher, T.S., et al. 1987. Primary structure of human pancreatic elastase 2 determined by sequence analysis of the cloned mRNA. *Biochemistry* 26: 7256-7261.
2. Kawashima, I., et al. 1987. Characterization of pancreatic elastase II cDNAs: two elastase II mRNAs are expressed in human pancreas. *DNA* 6: 163-172.
3. Moulard, M., et al. 1990. Further studies on the human pancreatic binary complexes involving procarboxypeptidase A. *FEBS Lett.* 261: 179-183.
4. Online Mendelian Inheritance in Man, OMIM<sup>™</sup>. 2005. Johns Hopkins University, Baltimore, MD. MIM Number: 609443. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
5. Szepessy, E., et al. 2006. Inactivity of recombinant ELA2B provides a new example of evolutionary elastase silencing in humans. *Pancreatology* 6: 117-122.

## CHROMOSOMAL LOCATION

Genetic locus: CELA2B (human) mapping to 1p36.21.

## PRODUCT

Elastase-2B siRNA (h) is a pool of 2 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Elastase-2B shRNA Plasmid (h): sc-105327-SH and Elastase-2B shRNA (h) Lentiviral Particles: sc-105327-V as alternate gene silencing products.

For independent verification of Elastase-2B (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-105327A and sc-105327B.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

Elastase-2B siRNA (h) is recommended for the inhibition of Elastase-2B expression in human cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Elastase-2B gene expression knockdown using RT-PCR Primer: Elastase-2B (h)-PR: sc-105327-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.