

FMO4 siRNA (m): sc-105365

BACKGROUND

The flavin-containing monooxygenase (FMO) family consists of five gene products, FMO1-5, that are major enzymatic oxidants involved in the metabolism of various therapeutics. Amino-trimethylamine (TMA), a diet-derived chemical from eggs, fish and legumes, is metabolized by FMOs. A polymorphism in genes encoding FMOs leads to a reduced TMA amino-oxidation capacity, leading to the excretion of relatively large amounts of TMA in urine, sweat and breath. This condition is known as trimethylaminuria, also known as fish-odor syndrome because individuals with this polymorphism exhibit a fishy body odor due to the free, unmetabolized amine. Located in the liver, FMO4 (flavin-containing monooxygenase 4), also known as Dimethylaniline monooxygenase and originally termed FMO2, is a 558 amino acid endoplasmic reticular protein that shares about 50-percent sequence similarity with FMO1.

REFERENCES

1. Dolphin, C.T., Shephard, E.A., Povey, S., Smith, R.L. and Phillips, I.R. 1992. Cloning, primary sequence and chromosomal localization of human FMO2, a new member of the flavin-containing mono-oxygenase family. *Biochem. J.* 287: 261-267.
2. Lawton, M.P., Cashman, J.R., Cresteil, T., Dolphin, C.T., Elfarra, A.A., Hines, R.N., Hodgson, E., Kimura, T., Ozols, J. and Phillips, I.R. 1994. A nomenclature for the mammalian flavin-containing monooxygenase gene family based on amino acid sequence identities. *Arch. Biochem. Biophys.* 308: 254-257.
3. Phillips, I.R., Dolphin, C.T., Clair, P., Hadley, M.R., Hutt, A.J., McCombie, R.R., Smith, R.L. and Shephard, E.A. 1995. The molecular biology of the flavin-containing monooxygenases of man. *Chem. Biol. Interact.* 96: 17-32.
4. Furnes, B., Feng, J., Sommer, S.S. and Schlenk, D. 2003. Identification of novel variants of the flavin-containing monooxygenase gene family in African Americans. *Drug Metab. Dispos.* 31: 187-193.
5. Cashman, J.R. 2008. Role of flavin-containing monooxygenase in drug development. *Expert Opin. Drug Metab. Toxicol.* 4: 1507-1521.
6. Henderson, M.C., Siddens, L.K., Morre, J.T., Krueger, S.K. and Williams, D.E. 2008. Metabolism of the anti-tuberculosis drug ethionamide by mouse and human FMO1, FMO2 and FMO3 and mouse and human lung microsomes. *Toxicol. Appl. Pharmacol.* 233: 420-427.
7. Online Mendelian Inheritance in Man, OMIM[™]. 2008. Johns Hopkins University, Baltimore, MD. MIM Number: 136131. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
8. Zhang, J., Chaluvadi, M.R., Reddy, R., Motika, M.S., Richardson, T.A., Cashman, J.R. and Morgan, E.T. 2009. Hepatic flavin-containing monooxygenase gene regulation in different mouse inflammation models. *Drug Metab. Dispos.* 37: 462-468.
9. Novick, R.M., Mitzey, A.M., Brownfield, M.S. and Elfarra, A.A. 2009. Differential localization of flavin-containing monooxygenase (FMO) isoforms 1, 3, and 4 in rat liver and kidney and evidence for expression of FMO4 in mouse, rat, and human liver and kidney microsomes. *J. Pharmacol. Exp. Ther.* 329: 1148-1155.

CHROMOSOMAL LOCATION

Genetic locus: Fmo4 (mouse) mapping to 1 H2.1.

PRODUCT

FMO4 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see FMO4 shRNA Plasmid (m): sc-105365-SH and FMO4 shRNA (m) Lentiviral Particles: sc-105365-V as alternate gene silencing products.

For independent verification of FMO4 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-105365A, sc-105365B and sc-105365C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

FMO4 siRNA (m) is recommended for the inhibition of FMO4 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor FMO4 gene expression knockdown using RT-PCR Primer: FMO4 (m)-PR: sc-105365-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.