

GPR52 siRNA (m): sc-105409

BACKGROUND

G protein-coupled receptors (GPRs), also known as seven transmembrane receptors, heptahelical receptors or 7TM receptors, comprise a superfamily of proteins that play a role in many different stimulus-response pathways. G protein-coupled receptors translate extracellular signals into intracellular signals (G protein activation) and they respond to a variety of signaling molecules, such as hormones and neurotransmitters. GPR52 (G protein-coupled receptor 52) is a 361 amino acid multi-pass membrane protein that belongs to the G protein-coupled receptor 1 family and has been found to be expressed in the caudate and putamen regions of the brain. The gene encoding GPR52 maps to human chromosome 1, the largest human chromosome spanning about 260 million base pairs and making up 8% of the human genome.

REFERENCES

1. Larhammar, D., Blomqvist, A.G. and Wahlestedt, C. 1993. The receptor revolution—multiplicity of G protein-coupled receptors. *Drug Des. Discov.* 9: 179-188.
2. Ji, T.H., Grossmann, M. and Ji, I. 1998. G protein-coupled receptors. I. Diversity of receptor-ligand interactions. *J. Biol. Chem.* 273: 17299-17302.
3. Sawzdargo, M., Nguyen, T., Lee, D.K., Lynch, K.R., Cheng, R., Heng, H.H., George, S.R. and O'Dowd, B.F. 1999. Identification and cloning of three novel human G protein-coupled receptor genes GPR52, PsiGPR53 and GPR55: GPR55 is extensively expressed in human brain. *Brain Res. Mol. Brain Res.* 64: 193-198.
4. Schöneberg, T., Schultz, G. and Gudermann, T. 1999. Structural basis of G protein-coupled receptor function. *Mol. Cell. Endocrinol.* 151: 181-193.
5. Lee, D.K., Nguyen, T., Lynch, K.R., Cheng, R., Vanti, W.B., Arkhitko, O., Lewis, T., Evans, J.F., George, S.R. and O'Dowd, B.F. 2001. Discovery and mapping of ten novel G protein-coupled receptor genes. *Gene* 275: 83-91.
6. Wittenberger, T., Schaller, H.C. and Hellebrand, S. 2001. An expressed sequence tag (EST) data mining strategy succeeding in the discovery of new G protein-coupled receptors. *J. Mol. Biol.* 307: 799-813.
7. Online Mendelian Inheritance in Man, OMIM[™]. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 604106. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: Gpr52 (mouse) mapping to 1 H2.1.

PRODUCT

GPR52 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see GPR52 shRNA Plasmid (m): sc-105409-SH and GPR52 shRNA (m) Lentiviral Particles: sc-105409-V as alternate gene silencing products.

For independent verification of GPR52 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-105409A, sc-105409B and sc-105409C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

GPR52 siRNA (m) is recommended for the inhibition of GPR52 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor GPR52 gene expression knockdown using RT-PCR Primer: GPR52 (m)-PR: sc-105409-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.