

group IID sPLA₂ siRNA (m): sc-105417

BACKGROUND

Phospholipase A₂s (PLA₂s) constitute a family of esterases that hydrolyze the sn-2-acyl ester bond in glycerophospholipid molecules. These enzymes are generally calcium-dependent and have been found both intra- and extracellularly. By hydrolyzing the sn-2 bond in glycerophospholipids, PLA₂s release fatty acids. One such fatty acid, arachidonic acid, generates the substrates for the initiation of the arachidonic acid cascade that produces various eicosanoids, many of which are potent mediators of inflammation. As a member of the PLA₂ family, group IID sPLA₂, also known as PLA2G2D (phosphatidylcholine 2-acylhydrolase 2D), sPLA₂-IID or SPLASH, is a 145 amino acid secreted protein that is broadly expressed but found at highest levels in pancreas and spleen. Group IID sPLA₂ catalyzes the reaction of phosphatidylcholine and water into 1-acylglycerophosphocholine and carboxylate, and has been linked to body weight loss in patients with chronic obstructive pulmonary disease (COPD), as well as lymphotoxin deficiency.

REFERENCES

1. Ishizaki, J., Suzuki, N., Higashino, K., Yokota, Y., Ono, T., Kawamoto, K., Fujii, N., Arita, H. and Hanasaki, K. 1999. Cloning and characterization of novel mouse and human secretory phospholipase A₂s. *J. Biol. Chem.* 274: 24973-24979.
2. Shakhov, A.N., Rubtsov, A.V., Lyakhov, I.G., Tumanov, A.V. and Nedospasov, S.A. 2000. SPLASH (PLA₂-IID), a novel member of phospholipase A₂ family, is associated with lymphotoxin deficiency. *Genes Immun.* 1: 191-199.
3. Online Mendelian Inheritance in Man, OMIM[™]. 2001. Johns Hopkins University, Baltimore, MD. MIM Number: 605630. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
4. Ecker, R., Menschikowski, M., Lattke, P. and Jaross, W. 2004. Increased hepatic cholesterol accumulation in transgenic mice overexpressing human secretory phospholipase A₂ group IIA. *Inflammation* 28: 59-65.
5. Takabatake, N., Sata, M., Inoue, S., Shibata, Y., Abe, S., Wada, T., Machiya, J., Ji, G., Matsuura, T., Takeishi, Y., Muramatsu, M. and Kubota, I. 2005. A novel polymorphism in secretory phospholipase A₂-IID is associated with body weight loss in chronic obstructive pulmonary disease. *Am. J. Respir. Crit. Care Med.* 172: 1097-1104.
6. Lindbom, J., Ljungman, A.G. and Tagesson, C. 2005. Interferon γ -induced gene expression of the novel secretory phospholipase A₂ type IID in human monocyte-derived macrophages is inhibited by lipopolysaccharide. *Inflammation* 29: 108-117.
7. Igarashi, A., Shibata, Y., Yamauchi, K., Osaka, D., Takabatake, N., Abe, S., Inoue, S., Kimura, T., Yamaguchi, Y., Ishizaki, J., Hanasaki, K. and Kubota, I. 2009. Gly80Ser polymorphism of phospholipase A₂-IID is associated with cytokine inducibility in A549 cells. *Respiration* 78: 312-321.

CHROMOSOMAL LOCATION

Genetic locus: Pla2g2d (mouse) mapping to 4 D3.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

group IID sPLA₂ siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see group IID sPLA₂ shRNA Plasmid (m): sc-105417-SH and group IID sPLA₂ shRNA (m) Lentiviral Particles: sc-105417-V as alternate gene silencing products.

For independent verification of group IID sPLA₂ (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-105417A, sc-105417B and sc-105417C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

group IID sPLA₂ siRNA (m) is recommended for the inhibition of group IID sPLA₂ expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor group IID sPLA₂ gene expression knockdown using RT-PCR Primer: group IID sPLA₂ (m)-PR: sc-105417-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.