



Importin-9 siRNA (m): sc-105575

BACKGROUND

The Importin complex consists of Importin- α and Importin- β proteins which assist in transport of arginine- or serine-rich (SR) proteins across the nucleus. Importin-9, also known as Imp9 or IPO9, is a 1,041 amino acid protein that contains one importin N-terminal domain and belongs to the Importin- β family. Localized to both the nucleus and the cytoplasm, Importin-9 functions as a nuclear transport receptor that mediates the docking of the Importin complex to the nuclear pore complex (NPC). Importin-9 mediates nuclear import of H2B histone, Ribosomal Protein S7 and Ribosomal Protein L18A. The gene encoding Importin-9 is located on human chromosome 1, the largest human chromosome that spans about 260 million base pairs and makes up 8% of the human genome.

REFERENCES

1. Watson, M.L., et al. 1990. Genomic organization of the selectin family of leukocyte adhesion molecules on human and mouse chromosome 1. *J. Exp. Med.* 172: 263-272.
2. Mühlgässer, P., et al. 2001. Multiple pathways contribute to nuclear import of core histones. *EMBO Rep.* 2: 690-696.
3. Jäkel, S., et al. 2002. Importins fulfil a dual function as nuclear import receptors and cytoplasmic chaperones for exposed basic domains. *EMBO J.* 21: 377-386.
4. Lubert, E.J., et al. 2003. Interaction between protein phosphatase 2A and members of the importin β superfamily. *Biochem. Biophys. Res. Commun.* 303: 908-913.
5. Kortvely, E., et al. 2005. Cloning and characterization of rat importin 9: implication for its neuronal function. *Brain Res. Mol. Brain Res.* 139: 103-114.
6. Weise, A., et al. 2005. New insights into the evolution of chromosome 1. *Cytogenet. Genome Res.* 108: 217-222.
7. Gregory, S.G., et al. 2006. The DNA sequence and biological annotation of human chromosome 1. *Nature* 441: 315-321.
8. Waldmann, I., et al. 2007. Nuclear import of c-Jun is mediated by multiple transport receptors. *J. Biol. Chem.* 282: 27685-27692.

CHROMOSOMAL LOCATION

Genetic locus: Ipo9 (mouse) mapping to 1 E4.

PRODUCT

Importin-9 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Importin-9 shRNA Plasmid (m): sc-105575-SH and Importin-9 shRNA (m) Lentiviral Particles: sc-105575-V as alternate gene silencing products.

For independent verification of Importin-9 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-105575A, sc-105575B and sc-105575C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Importin-9 siRNA (m) is recommended for the inhibition of Importin-9 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Importin-9 gene expression knockdown using RT-PCR Primer: Importin-9 (m)-PR: sc-105575-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.