

ING5 siRNA (h): sc-105577

BACKGROUND

Inhibitor of growth protein (ING) family of nuclear proteins, also designated ING tumor suppressor proteins, inhibit tumor progression by modulating the transcriptional outputs of signaling pathways, which in turn regulates cell proliferation. Members of this family include ING1, ING2, ING3, ING4, ING5 and INGX. ING5 (inhibitor of growth family, member 5), also known as p28ING5, is a 240 amino acid nuclear protein containing a single PHD-type zinc finger, a common motif found in proteins involved in chromatin remodeling. Through chromatin acetylation, ING5 may regulate DNA replication and may function as a transcriptional coactivator. ING5 also acts as a tumor suppressor that interacts with p53, inhibits cell growth and induces apoptosis. ING5 is expressed as two isoforms produced by alternative splicing events and is encoded by a gene located on human chromosome 2.

REFERENCES

- Shiseki, M., Nagashima, M., Pedoux, R.M., Kitahama-Shiseki, M., Miura, K., Okamura, S., Onogi, H., Higashimoto, Y., Appella, E., Yokota, J. and Harris, C.C. 2003. p29ING4 and p28ING5 bind to p53 and p300, and enhance p53 activity. *Cancer Res.* 63: 2373-2378.
- Gong, W., Suzuki, K., Russell, M. and Riabowol, K. 2005. Function of the ING family of PHD proteins in cancer. *Int. J. Biochem. Cell Biol.* 37: 1054-1065.
- He, G.H., Helbing, C.C., Wagner, M.J., Sensen, C.W. and Riabowol, K. 2005. Phylogenetic analysis of the ING family of PHD finger proteins. *Mol. Biol. Evol.* 22: 104-116.
- Doyon, Y., Cayrou, C., Ullah, M., Landry, A.J., Côte, V., Selleck, W., Lane, W.S., Tan, S., Yang, X.J. and Côte, J. 2006. ING tumor suppressor proteins are critical regulators of chromatin acetylation required for genome expression and perpetuation. *Mol. Cell* 21: 51-64.

CHROMOSOMAL LOCATION

Genetic locus: ING5 (human) mapping to 2q37.3.

PRODUCT

ING5 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ING5 shRNA Plasmid (h): sc-105577-SH and ING5 shRNA (h) Lentiviral Particles: sc-105577-V as alternate gene silencing products.

For independent verification of ING5 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-105577A, sc-105577B and sc-105577C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

ING5 siRNA (h) is recommended for the inhibition of ING5 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

ING5 (36.85): sc-135745 is recommended as a control antibody for monitoring of ING5 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-mouse IgG-HRP: sc-2005 (dilution range: 1:2000-1:32,000) or Cruz Marker™ compatible goat anti-mouse IgG-HRP: sc-2031 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use goat anti-mouse IgG-FITC: sc-2010 (dilution range: 1:100-1:400) or goat anti-mouse IgG-TR: sc-2781 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ING5 gene expression knockdown using RT-PCR Primer: ING5 (h)-PR: sc-105577-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.