

T2-cadherin siRNA (h): sc-106589

BACKGROUND

The cadherins are a family of Ca^{2+} -dependent adhesion molecules that function to mediate cell-cell binding critical to the maintenance of structure and morphogenesis. Cadherins each contain a large extracellular domain at the N-terminus, which is characterized by a series of five homologous repeats, the most distal of which is thought to be responsible for binding specificity. T2-cadherin, also known as cadherin-10 or CDH10, is a 788 amino acid single-pass type I membrane protein that belongs to the cadherin family and contains five cadherin domains. Expressed predominately in brain, but also found in prostate and both adult and fetal kidney, T2-cadherin functions as a calcium-dependent cell adhesion protein that is thought to be involved in the maintenance of synaptic adhesions, axon outgrowth and guidance.

REFERENCES

1. Suzuki, S., Sano, K. and Tanihara, H. 1991. Diversity of the cadherin family: evidence for eight new cadherins in nervous tissue. *Cell Regul.* 2: 261-270.
2. Kremmidiotis, G., Baker, E., Crawford, J., Eyre, H.J., Nahmias, J. and Callen, D.F. 1998. Localization of human cadherin genes to chromosome regions exhibiting cancer-related loss of heterozygosity. *Genomics* 49: 467-471.
3. Kools, P., Vanhalst, K., Van den Eynde, E. and van Roy, F. 1999. The human cadherin-10 gene: complete coding sequence, predominant expression in the brain, and mapping on chromosome 5p13-14. *FEBS Lett.* 452: 328-334.
4. Shimoyama, Y., Tsujimoto, G., Kitajima, M. and Natori, M. 2000. Identification of three human type-II classic cadherins and frequent heterophilic interactions between different subclasses of type-II classic cadherins. *Biochem. J.* 349: 159-167.
5. Online Mendelian Inheritance in Man, OMIM[™]. 2000. Johns Hopkins University, Baltimore, MD. MIM Number: 604555. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
6. Gil, O.D., Needleman, L. and Huntley, G.W. 2002. Developmental patterns of cadherin expression and localization in relation to compartmentalized thalamocortical terminations in rat barrel cortex. *J. Comp. Neurol.* 453: 372-388.
7. Williams, M.J., Lowrie, M.B., Bennett, J.P., Firth, J.A. and Clark, P. 2005. Cadherin-10 is a novel blood-brain barrier adhesion molecule in human and mouse. *Brain Res.* 1058: 62-72.
8. Walker, M.M., Ellis, S.M., Auza, M.J., Patel, A. and Clark, P. 2008. The intercellular adhesion molecule, cadherin-10, is a marker for human prostate luminal epithelial cells that is not expressed in prostate cancer. *Mod. Pathol.* 21: 85-95.

CHROMOSOMAL LOCATION

Genetic locus: CDH10 (human) mapping to 5p14.2.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

T2-cadherin siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see T2-cadherin shRNA Plasmid (h): sc-106589-SH and T2-cadherin shRNA (h) Lentiviral Particles: sc-106589-V as alternate gene silencing products.

For independent verification of T2-cadherin (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-106589A, sc-106589B and sc-106589C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20°C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20°C , avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μl of RNase-free water makes a 10 μM solution in a 10 μM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

T2-cadherin siRNA (h) is recommended for the inhibition of T2-cadherin expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μM in 66 μl . Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor T2-cadherin gene expression knockdown using RT-PCR Primer: T2-cadherin (h)-PR: sc-106589-PR (20 μl). Annealing temperature for the primers should be $55-60^{\circ}\text{C}$ and the extension temperature should be $68-72^{\circ}\text{C}$.

RESEARCH USE

For research use only, not for use in diagnostic procedures.