

USP50 siRNA (m): sc-106682

BACKGROUND

The ubiquitin (Ub) pathway involves three sequential enzymatic steps that facilitate the conjugation of Ub and Ub-like molecules to specific protein substrates. Through the use of a wide range of enzymes that can add or remove ubiquitin, the Ub pathway controls many intracellular processes such as signal transduction, transcriptional activation and cell cycle progression. USP50 (ubiquitin specific peptidase 50) is a 339 amino acid protein that belongs to the peptidase C19 family of proteins. Weakly expressed in a few tissues, USP50 contains all of the active residues necessary to function as a deubiquitinating enzyme, but it appears to be catalytically inactive. The gene encoding USP50 maps to human chromosome 15, which houses over 700 genes and comprises nearly 3% of the human genome. Angelman syndrome, Prader-Willi syndrome, Tay-Sachs disease and Marfan syndrome are all associated with defects in chromosome 15-localized genes.

REFERENCES

1. Puente, X.S., Sánchez, L.M., Overall, C.M. and López-Otín, C. 2003. Human and mouse proteases: a comparative genomic approach. *Nat. Rev. Genet.* 4: 544-558.
2. Quesada, V., Díaz-Perales, A., Gutiérrez-Fernández, A., Garabaya, C., Cal, S. and López-Otín, C. 2004. Cloning and enzymatic analysis of 22 novel human ubiquitin-specific proteases. *Biochem. Biophys. Res. Commun.* 314: 54-62.
3. Hicke, L., Schubert, H.L. and Hill, C.P. 2005. Ubiquitin-binding domains. *Nat. Rev. Mol. Cell Biol.* 6: 610-621.
4. Espeseth, A.S., Huang, Q., Gates, A., Xu, M., Yu, Y., Simon, A.J., Shi, X.P., Zhang, X., Hodor, P., Stone, D.J., Burchard, J., Cavet, G., Bartz, S., Linsley, P., Ray, W.J. and Hazuda, D. 2006. A genome wide analysis of ubiquitin ligases in APP processing identifies a novel regulator of BACE1 mRNA levels. *Mol. Cell. Neurosci.* 33: 227-235.
5. Stegmeier, F., Rape, M., Draviam, V.M., Nalepa, G., Sowa, M.E., Ang, X.L., McDonald, E.R., Li, M.Z., Hannon, G.J., Sorger, P.K., Kirschner, M.W., Harper, J.W. and Elledge, S.J. 2007. Anaphase initiation is regulated by antagonistic ubiquitination and deubiquitination activities. *Nature* 446: 876-881.
6. Ha, B.H. and Kim, E.E. 2008. Structures of proteases for ubiquitin and ubiquitin-like modifiers. *BMB Rep.* 41: 435-443.
7. Ventii, K.H. and Wilkinson, K.D. 2008. Protein partners of deubiquitinating enzymes. *Biochem. J.* 414: 161-175.
8. Vos, R.M., Altreuter, J., White, E.A. and Howley, P.M. 2009. The ubiquitin-specific peptidase USP15 regulates human papillomavirus type 16 E6 protein stability. *J. Virol.* 83: 8885-8892.
9. Kirkin, V., McEwan, D.G., Novak, I. and Dikic, I. 2009. A role for ubiquitin in selective autophagy. *Mol. Cell* 34: 259-269.

CHROMOSOMAL LOCATION

Genetic locus: Usp50 (mouse) mapping to 2 F1.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

USP50 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see USP50 shRNA Plasmid (m): sc-106682-SH and USP50 shRNA (m) Lentiviral Particles: sc-106682-V as alternate gene silencing products.

For independent verification of USP50 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-106682A, sc-106682B and sc-106682C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

USP50 siRNA (m) is recommended for the inhibition of USP50 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor USP50 gene expression knockdown using RT-PCR Primer: USP50 (m)-PR: sc-106682-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.