

DGK- η siRNA (h): sc-106793

BACKGROUND

Diacylglycerol kinases (DGKs) phosphorylate diacylglycerol (DAG) to produce phosphatidic acid. DAG and phosphatidic acid are lipids that act as second messengers in signaling cascades. DGK- α influences cell activation and secretion of lethal exosomes, which in turn control cell death. DGK- β is abundant in restricted brain regions such as the caudate putamen and olfactory tubercle. DGK- γ encodes full-length and truncated transcripts that are present in a range of human tissues, with greatest expression observed in retina. DGK- δ is most abundant in skeletal muscle. DGK- ϵ shows specificity for arachidonyl-containing diacylglycerol and is expressed predominantly in testis. DGK- ζ is most abundant in brain and muscle. DGK- η is closely related to DGK- δ and contains one PH domain, two phorbol-ester/DAG-type zinc fingers and one SAM (sterile alpha motif) domain. DGK- θ is most abundant in the cerebellum and hippocampus. DGK- ι is present in brain and retina as a predominant transcript of more than 12 kb, including a long 3' untranslated region, with additional low abundance transcripts of 9.5 and 7.5 kb.

REFERENCES

1. Schaap, D., et al. 1990. Purification, cDNA-cloning and expression of human diacylglycerol kinase. *FEBS Lett.* 275: 151-158.
2. Goto, K., et al. 1993. Molecular cloning and expression of a 90-kDa diacylglycerol kinase that predominantly localizes in neurons. *Proc. Natl. Acad. Sci. USA* 90: 7598-7602.
3. Masai, I., et al. 1993. *Drosophila* retinal degeneration A gene encodes an eye-specific diacylglycerol kinase with cysteine-rich zinc-finger motifs and ankyrin repeats. *Proc. Natl. Acad. Sci. USA* 90: 11157-11161.
4. Kai, M., et al. 1994. Molecular cloning of a diacylglycerol kinase isozyme predominantly expressed in human retina with a truncated and inactive enzyme expression in most other human cells. *J. Biol. Chem.* 269: 18492-18498.
5. Sakane, F., et al. 1996. Molecular cloning of a novel diacylglycerol kinase isozyme with a pleckstrin homology domain and a C-terminal tail similar to those of the EPH family of protein-tyrosine kinases. *J. Biol. Chem.* 271: 8394-8401.
6. Tang, W., et al. 1996. Molecular cloning of a novel human diacylglycerol kinase highly selective for arachidonate-containing substrates. *J. Biol. Chem.* 271: 10237-10241.
7. Klauk, T.M., et al. 1996. Cloning and characterization of a glucocorticoid-induced diacylglycerol kinase. *J. Biol. Chem.* 271: 19781-19788.
8. Ding, L., et al. 1998. The cloning and characterization of a novel human diacylglycerol kinase, DGK ι . *J. Biol. Chem.* 273: 32746-32752.
9. Topham, M.K., et al. 1999. Mammalian diacylglycerol kinases, a family of lipid kinases with signaling functions. *J. Biol. Chem.* 274: 11447-11450.

CHROMOSOMAL LOCATION

Genetic locus: DGKH (human) mapping to 13q14.11.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

DGK- η siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see DGK- η shRNA Plasmid (h): sc-106793-SH and DGK- η shRNA (h) Lentiviral Particles: sc-106793-V as alternate gene silencing products.

For independent verification of DGK- η (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-106793A, sc-106793B and sc-106793C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

DGK- η siRNA (h) is recommended for the inhibition of DGK- η expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor DGK- η gene expression knockdown using RT-PCR Primer: DGK- η (h)-PR: sc-106793-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.