

PCDHGA8 siRNA (h): sc-106828

BACKGROUND

Protocadherins are a large family of cadherin-like cell adhesion proteins that are involved in the establishment and maintenance of neuronal connections in the brain. There are three protocadherin gene clusters designated α , β and γ , all of which contain multiple tandemly arranged genes. The protocadherin γ cluster consists of three subfamilies (A, B and C). As a member of the γ subfamily A, PCDHGA8 (protocadherin γ A8) is a 932 amino acid protein that is one of 22 proteins encoded by the protocadherin γ cluster. Typical of γ protocadherins, PCDHGA8 contains six cadherin motifs and is a type I transmembrane receptor expressed in the central nervous system. With localization to synapses, members of the γ cluster of protocadherins are essential for neuronal survival. There are two isoforms of PCDHGA8 that are produced as a result of alternative splicing events.

REFERENCES

1. Wu, Q. and Maniatis, T. 1999. A striking organization of a large family of human neural cadherin-like cell adhesion genes. *Cell* 97: 779-790.
2. Wu, Q., et al. 2001. Comparative DNA sequence analysis of mouse and human protocadherin gene clusters. *Genome Res.* 11: 389-404.
3. Wang, X., et al. 2002. γ protocadherins are required for survival of spinal interneurons. *Neuron* 36: 843-854.
4. Kirov, G., et al. 2003. Variation in the protocadherin γ A gene cluster. *Genomics* 82: 433-440.
5. Frank, M., et al. 2005. Differential expression of individual γ -protocadherins during mouse brain development. *Mol. Cell. Neurosci.* 29: 603-616.
6. Reiss, K., et al. 2006. Regulated ADAM10-dependent ectodomain shedding of γ -protocadherin C3 modulates cell-cell adhesion. *J. Biol. Chem.* 281: 21735-21744.
7. Bonn, S., et al. 2007. Combinatorial expression of α - and γ -protocadherins alters their presenilin-dependent processing. *Mol. Cell. Biol.* 27: 4121-4132.
8. Online Mendelian Inheritance in Man, OMIM[™]. 2007. Johns Hopkins University, Baltimore, MD. MIM Number: 606295. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: PCDHGA8 (human) mapping to 5q31.3.

PRODUCT

PCDHGA8 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see PCDHGA8 shRNA Plasmid (h): sc-106828-SH and PCDHGA8 shRNA (h) Lentiviral Particles: sc-106828-V as alternate gene silencing products.

For independent verification of PCDHGA8 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-106828A, sc-106828B and sc-106828C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

PCDHGA8 siRNA (h) is recommended for the inhibition of PCDHGA8 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

PCDHGA8 (NQ-B21): sc-134418 is recommended as a control antibody for monitoring of PCDHGA8 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor PCDHGA8 gene expression knockdown using RT-PCR Primer: PCDHGA8 (h)-PR: sc-106828-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.