

# ALG-2 siRNA (h): sc-106841

## BACKGROUND

An increased intracellular  $\text{Ca}^{2+}$  concentration induces apoptotic cell death. Transiently elevated  $\text{Ca}^{2+}$  concentrations are required for glucocorticoid-mediated and T cell receptor-mediated pathways, leading to T cell apoptosis. ALG-2 (for apoptosis-linked gene 2) is a  $\text{Ca}^{2+}$ -binding protein that participates in regulatory events occurring late in the apoptotic program and where several death signals converge. ALG-2 is a protein expressed in normal brain, and to a greater extent in ischemic brain. The ALG-2 protein contains five EF-hand-like motifs and shares homology with members of the penta EF-hand family, which includes Calpain small subunits sorcin and Grancalcin.

## REFERENCES

1. McConkey, D.J., et al. 1989. Calcium-dependent killing of immature thymocytes by stimulation via the CD3/T cell receptor complex. *J. Immunol.* 143: 1801-1806.
2. McConkey, D.J., et al. 1989. Glucocorticoids activate a suicide process in thymocytes through an elevation of cytosolic  $\text{Ca}^{2+}$  concentration. *Arch. Biochem. Biophys.* 269: 365-370.
3. Nicotera, P., et al. 1990. The role of  $\text{Ca}^{2+}$  in cell killing. *Chem. Res. Toxicol.* 3: 484-494.
4. Vito, P., et al. 1996. Interfering with apoptosis:  $\text{Ca}^{2+}$ -binding protein ALG-2 and Alzheimer's disease gene ALG-3. *Science* 271: 521-525.
5. D'Adamio, L., et al. 1997. Functional cloning of genes involved in T cell receptor-induced programmed cell death. *Semin. Immunol.* 9: 17-23.
6. Maki, M., et al. 1997. A growing family of the  $\text{Ca}^{2+}$ -binding proteins with five EF-hand motifs. *Biochem. J.* 328: 718-720.

## CHROMOSOMAL LOCATION

Genetic locus: PDC6 (human) mapping to 5p15.33.

## PRODUCT

ALG-2 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu\text{M}$  solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ALG-2 shRNA Plasmid (h): sc-106841-SH and ALG-2 shRNA (h) Lentiviral Particles: sc-106841-V as alternate gene silencing products.

For independent verification of ALG-2 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-106841A, sc-106841B and sc-106841C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at  $-20^{\circ}\text{C}$  with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at  $-20^{\circ}\text{C}$ , avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu\text{l}$  of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu\text{l}$  of RNase-free water makes a 10  $\mu\text{M}$  solution in a 10  $\mu\text{M}$  Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

ALG-2 siRNA (h) is recommended for the inhibition of ALG-2 expression in human cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu\text{M}$  in 66  $\mu\text{l}$ . Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## GENE EXPRESSION MONITORING

ALG-2 (H-11): sc-376950 is recommended as a control antibody for monitoring of ALG-2 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG $\kappa$  BP-FITC: sc-516140 or m-IgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ALG-2 gene expression knockdown using RT-PCR Primer: ALG-2 (h)-PR: sc-106841-PR (20  $\mu\text{l}$ ). Annealing temperature for the primers should be  $55-60^{\circ}\text{C}$  and the extension temperature should be  $68-72^{\circ}\text{C}$ .

## SELECT PRODUCT CITATIONS

1. Cho, H.J. and Mook-Jung, I. 2018. O-GlcNAcylation regulates endoplasmic reticulum exit sites through Sec31A modification in conventional secretory pathway. *FASEB J.* 32: 4641-4657.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.