

NOMO3 siRNA (h): sc-106857

BACKGROUND

Three highly similar proteins termed NOMO1, NOMO2 and NOMO3, are encoded by a gene mapping to a region of duplication on the p arm of human chromosome 16. All three NOMO proteins share similar functions and have been difficult to characterize individually. NOMO1 (Nodal modulator 1), also known as PM5, is a 1,222 amino acid highly conserved single-pass type I membrane protein expressed in colon tumor tissue and normal colonic mucosa. NOMO proteins are novel antagonists of Nodal signaling which interact with Nicalin to form a Nicalin-NOMO complex, and are rapidly degraded or stabilized by Nicalin. NOMO proteins were once considered candidates for the development of pseudoxanthoma elasticum (PXE), a heritable disorder of connective tissue, as the NOMO genes are located in close proximity to the gene responsible for PXE development (MRP6).

REFERENCES

1. Templeton, N.S., Rodgers, L.A., Levy, A.T., Ting, K.L., Krutzsch, H.C., Liotta, L.A. and Stetler-Stevenson, W.G. 1992. Cloning and characterization of a novel human cDNA that has DNA similarity to the conserved region of the collagenase gene family. *Genomics* 12: 175-176.
2. Loftus, B.J., Kim, U.J., Sneddon, V.P., Kalush, F., Brandon, R., Fuhrmann, J., Mason, T., Crosby, M.L., Barnstead, M., Cronin, L., Deslattes Mays, A., Cao, Y., Xu, R.X., Kang, H.L., Mitchell, S., Eichler, E.E., Harris, P.C., et al. 1999. Genome duplications and other features in 12 Mb of DNA sequence from human chromosome 16p and 16q. *Genomics* 60: 295-308.
3. Perdu, J. and Germain, D.P. 2001. Identification of novel polymorphisms in the pM5 and MRP1 (ABCC1) genes at locus 16p13.1 and exclusion of both genes as responsible for pseudoxanthoma elasticum. *Hum. Mutat.* 17: 74-75.
4. Haffner, C., Frauli, M., Topp, S., Irmeler, M., Hofmann, K., Regula, J.T., Bally-Cuif, L. and Haass, C. 2004. Nicalin and its binding partner Nomo are novel Nodal signaling antagonists. *EMBO J.* 23: 3041-3050.
5. Online Mendelian Inheritance in Man, OMIM[™]. 2005. Johns Hopkins University, Baltimore, MD. MIM Number: 609157. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
6. Haffner, C. and Haass, C. 2006. Cellular functions of γ -secretase-related proteins. *Neurodegener. Dis.* 3: 284-289.
7. Haffner, C., Dettmer, U., Weiler, T. and Haass, C. 2007. The Nicastrin-like protein Nicalin regulates assembly and stability of the Nicalin-Nodal modulator (NOMO) membrane protein complex. *J. Biol. Chem.* 282: 10632-10638.

CHROMOSOMAL LOCATION

Genetic locus: NOMO3 (human) mapping to 16p13.11.

PRODUCT

NOMO3 siRNA (h) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see NOMO3 shRNA Plasmid (h): sc-106857-SH and NOMO3 shRNA (h) Lentiviral Particles: sc-106857-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

NOMO3 siRNA (h) is recommended for the inhibition of NOMO3 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor NOMO3 gene expression knockdown using RT-PCR Primer: NOMO3 (h)-PR: sc-106857-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.