

PCDHGC5 siRNA (h): sc-106934

BACKGROUND

The PCDHGA6 gene is a member of the protocadherin γ gene cluster, one of three related clusters tandemly linked on chromosome 5q31.3. These gene clusters have an immunoglobulin-like organization, suggesting that a novel mechanism may be involved in their regulation and expression. The γ gene cluster includes 22 genes divided into 3 subfamilies. Subfamily A contains 12 genes, subfamily B contains 7 genes and 2 pseudogenes, and the more distantly related subfamily C contains 3 genes. PCDHGC5 (protocadherin γ -C5) is a 944 amino acid single-pass type I membrane protein that contains six cadherin domains and exists as two alternatively spliced isoforms. As a potential calcium-dependent cell-adhesion protein, PCDHGC5 may be involved in the establishment and maintenance of specific neuronal connections in the brain. The gene that encodes PCDHGC5 consists of approximately 23,739 bases and maps to human chromosome 5q31.3.

REFERENCES

1. Wu, Q. and Maniatis, T. 1999. A striking organization of a large family of human neural cadherin-like cell adhesion genes. *Cell* 97: 779-790.
2. Yagi, T. and Takeichi, M. 2000. Cadherin superfamily genes: functions, genomic organization, and neurologic diversity. *Genes Dev.* 14: 1169-1180.
3. Nollet, F., Kools, P. and van Roy, F. 2000. Phylogenetic analysis of the cadherin superfamily allows identification of six major subfamilies besides several solitary members. *J. Mol. Biol.* 299: 551-572.
4. Wu, Q. and Maniatis, T. 2000. Large exons encoding multiple ectodomains are a characteristic feature of protocadherin genes. *Proc. Natl. Acad. Sci. USA* 97: 3124-3129.
5. Wu, Q., Zhang, T., Cheng, J.F., Kim, Y., Grimwood, J., Schmutz, J., Dickson, M., Noonan, J.P., Zhang, M.Q., Myers, R.M. and Maniatis, T. 2001. Comparative DNA sequence analysis of mouse and human protocadherin gene clusters. *Genome Res.* 11: 389-404.
6. Online Mendelian Inheritance in Man, OMIM[™]. 2001. Johns Hopkins University, Baltimore, MD. MIM Number: 606306. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
7. Abe, M., Watanabe, N., McDonnell, N., Takato, T., Ohira, M., Nakagawara, A. and Ushijima, T. 2008. Identification of genes targeted by CpG island methylator phenotype in neuroblastomas, and their possible integrative involvement in poor prognosis. *Oncology* 74: 50-60.
8. Wang, L., Srivastava, A.K. and Schwartz, C.E. 2010. Microarray data integration for genome-wide analysis of human tissue-selective gene expression. *BMC Genomics* 11: S15.

CHROMOSOMAL LOCATION

Genetic locus: PCDHGC5 (human) mapping to 5q31.3.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

PCDHGC5 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see PCDHGC5 shRNA Plasmid (h): sc-106934-SH and PCDHGC5 shRNA (h) Lentiviral Particles: sc-106934-V as alternate gene silencing products.

For independent verification of PCDHGC5 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-106934A, sc-106934B and sc-106934C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

PCDHGC5 siRNA (h) is recommended for the inhibition of PCDHGC5 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor PCDHGC5 gene expression knockdown using RT-PCR Primer: PCDHGC5 (h)-PR: sc-106934-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.