



TMEM248 siRNA (m): sc-108102

BACKGROUND

TMEM248 (transmembrane protein 248) is a 314 amino acid multi-pass membrane protein. Existing as two alternatively spliced isoforms, TMEM248 is encoded by a gene that maps to human chromosome 7q11.21 and mouse chromosome 5 G1.3. Chromosome 7 is about 158 million bases long, encodes over 1,000 genes and makes up about 5% of the human genome. Chromosome 7 has been linked to osteogenesis imperfecta, Pendred syndrome, lissencephaly, citrullinemia and Shwachman-Diamond syndrome. The deletion of a portion of the q arm of chromosome 7 is associated with Williams-Beuren syndrome, a condition characterized by mild mental retardation, an unusual comfort and friendliness with strangers and an elfin appearance. Deletions of portions of the q arm of chromosome 7 are also seen in a number of myeloid disorders including cases of acute myelogenous leukemia and myelodysplasia.

REFERENCES

1. Gupta, P., Rosen, J.M., D'Eustachio, P. and Ruddle, F.H. 1982. Localization of the casein gene family to a single mouse chromosome. *J. Cell Biol.* 93: 199-204.
2. Kozak, C.A. 1984. Differential expression of murine leukemia virus loci in chemically induced hybrid cells. *J. Virol.* 51: 876-879.
3. Cheng, S.V., Martin, G.R., Nadeau, J.H., Haines, J.L., Bucan, M., Kozak, C.A., MacDonald, M.E., Lockyer, J.L., Ledley, F.D. and Woo, S.L. 1989. Synteny on mouse chromosome 5 of homologs for human DNA loci linked to the Huntington disease gene. *Genomics* 4: 419-426.
4. McAllister, K.A., Ramachandran, S., Haugen-Strano, A., Fiedorek, F.T. and Wiseman, R.W. 1997. Genetic mapping of the BRCA2 breast cancer susceptibility gene on mouse chromosome 5. *Mamm. Genome* 8: 540-541.
5. Himmelbauer, H., Wedemeyer, N., Haaf, T., Wanker, E.E., Schalkwyk, L.C. and Lehrach, H. 1998. IRS-PCR-based genetic mapping of the Huntingtin interacting protein gene (HIP1) on mouse chromosome 5. *Mamm. Genome* 9: 26-31.
6. Shimizu, T. 1999. Mapping of a gene causing mouse gutter-shaped tooth root to chromosome 5. *Arch. Oral Biol.* 44: 917-924.

CHROMOSOMAL LOCATION

Genetic locus: Tmem248 (mouse) mapping to 5 G1.3.

PRODUCT

TMEM248 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TMEM248 shRNA Plasmid (m): sc-108102-SH and TMEM248 shRNA (m) Lentiviral Particles: sc-108102-V as alternate gene silencing products.

For independent verification of TMEM248 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-108102A, sc-108102B and sc-108102C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

TMEM248 siRNA (m) is recommended for the inhibition of TMEM248 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TMEM248 gene expression knockdown using RT-PCR Primer: TMEM248 (m)-PR: sc-108102-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.