

1110059E24Rik siRNA (m): sc-108191

BACKGROUND

1110059E24Rik is a 155 amino acid protein that is encoded by a gene which maps to mouse chromosome 19 B. Many of the mouse homologues of genes on chromosome 19 are located on human chromosome 10. Human chromosome 10 contains over 800 genes and 135 million nucleotides, making up nearly 4.5% of the human genome. PTEN is an important tumor suppressor gene located on chromosome 10 and, when defective, causes a genetic predisposition to cancer development known as Cowden syndrome. The chromosome 10 encoded gene ERCC6 is important for DNA repair and is linked to Cockayne syndrome which is characterized by extreme photosensitivity and premature aging. Tetrahydrobiopterin deficiency and a number of syndromes involving defective skull and facial bone fusion are also linked to chromosome 10. As with most trisomies, trisomy 10 is rare and is deleterious.

REFERENCES

1. Koch, G., Lalley, P.A., McAvoy, M. and Shows, T.B. 1981. Assignment of LIPA, associated with human acid lipase deficiency, to human chromosome 10 and comparative assignment to mouse chromosome 19. *Somatic Cell Genet.* 7: 345-358.
2. Yang-Feng, T.L., Landau, N.R., Baltimore, D. and Francke, U. 1986. The terminal deoxynucleotidyltransferase gene is located on human chromosome 10 (10q23—q24) and on mouse chromosome 19. *Cytogenet. Cell Genet.* 43: 121-126.
3. Mock, B.A., Connelly, M.A., McBride, O.W., Kozak, C.A. and Marcu, K.B. 1995. CHUK, a conserved helix-loop-helix ubiquitous kinase, maps to human chromosome 10 and mouse chromosome 19. *Genomics* 27: 348-351.
4. Luo, G., Leroy, E., Kozak, C.A., Polymeropoulos, M.H. and Horowitz, R. 1997. Mapping of the gene (NRAP) encoding N-RAP in the mouse and human genomes. *Genomics* 45: 229-232.
5. Deloukas, P., French, L., Meitinger, T. and Moschonas, N.K. 2000. Report of the third international workshop on human chromosome 10 mapping and sequencing 1999. *Cytogenet. Cell Genet.* 90: 1-12.
6. Freedman, B.I., Rich, S.S., Yu, H., Roh, B.H. and Bowden, D.W. 2002. Linkage heterogeneity of end-stage renal disease on human chromosome 10. *Kidney Int.* 62: 770-774.

CHROMOSOMAL LOCATION

Genetic locus: 1110059E24Rik (mouse) mapping to 19 B.

PRODUCT

1110059E24Rik siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see 1110059E24Rik shRNA Plasmid (m): sc-108191-SH and 1110059E24Rik shRNA (m) Lentiviral Particles: sc-108191-V as alternate gene silencing products.

For independent verification of 1110059E24Rik (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-108191A, sc-108191B and sc-108191C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

1110059E24Rik siRNA (m) is recommended for the inhibition of 1110059E24Rik expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor 1110059E24Rik gene expression knockdown using RT-PCR Primer: 1110059E24Rik (m)-PR: sc-108191-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.