

# β-1,4-Gal-T2 siRNA (m): sc-108224

## BACKGROUND

The β-1,4-Gal-T2 gene, which maps to chromosome 1p34.1 in human and 4 D2.1 in mouse, is one of seven β-1,4-galactosyltransferase (β-1,4-Gal-T) genes. These genes encode type II membrane-bound glycoproteins that appear to have exclusive specificity for the donor substrate UDP-galactose. These protein products transfer galactose in a β-1,4 linkage to similar acceptor sugars, such as GlcNAc, Glc and Xyl. These type II membrane glycoproteins have an N-terminal hydrophobic signal sequence that directs the protein to the Golgi apparatus and remains uncleaved to function as a transmembrane anchor. β-1,4-Gal-T2 (β-1,4-galactosyltransferase 2), also known as UDP-galactose:β-N-acetylglucosamine β-1,4-galactosyltransferase 2, is a 372 amino acid protein that is highly expressed in prostate, testis, ovary, intestine, muscle and fetal brain. β-1,4-Gal-T2 is responsible for the synthesis of complex-type N-linked oligosaccharides in many glycoproteins. Two isoforms of β-1,4-Gal-T2 exist as a result of alternative splicing events.

## REFERENCES

- Almeida, R., et al. 1997. A family of human β-4-galactosyltransferases. Cloning and expression of two novel UDP-galactose:β-n-acetylglucosamine β-1,4-galactosyltransferases, β-4Gal-T2 and β-4Gal-T3. *J. Biol. Chem.* 272: 31979-31991.
- Lo, N.W., et al. 1998. The expanding β-4-galactosyltransferase gene family: messages from the databanks. *Glycobiology* 8: 517-526.
- Amado, M., et al. 1999. Identification and characterization of large galactosyltransferase gene families: galactosyltransferases for all functions. *Biochim. Biophys. Acta* 1473: 35-53.
- Guo, S., et al. 2001. Galactosylation of N-linked oligosaccharides by human β-1,4-galactosyltransferases I, II, III, IV, V, and VI expressed in Sf-9 cells. *Glycobiology* 11: 813-820.
- Yan, M., et al. 2008. Expression change of β-1,4 galactosyltransferase I, V mRNAs and Galβ-1,4-GlcNAc group in rat sciatic nerve after crush. *J. Mol. Histol.* 39: 317-328.
- Kumagai, T., et al. 2009. Early lethality of β-1,4-galactosyltransferase V-mutant mice by growth retardation. *Biochem. Biophys. Res. Commun.* 379: 456-459.
- Mast, S.W., et al. 2009. Exploring the ultrastructural localization and biosynthesis of β-1,4-galactan in *Pinus radiata* compression wood. *Plant Physiol.* 150: 573-583.
- Yang, H., et al. 2009. Lipopolysaccharide induced upregulation of β-1,4-galactosyltransferase-I in Schwann cell. *Inflammation* 32: 279-286.

## CHROMOSOMAL LOCATION

Genetic locus: B4galt2 (mouse) mapping to 4 D2.1.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## PRODUCT

β-1,4-Gal-T2 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see β-1,4-Gal-T2 shRNA Plasmid (m): sc-108224-SH and β-1,4-Gal-T2 shRNA (m) Lentiviral Particles: sc-108224-V as alternate gene silencing products.

For independent verification of β-1,4-Gal-T2 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-108224A, sc-108224B and sc-108224C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μl of RNase-free water makes a 10 μM solution in a 10 μM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

β-1,4-Gal-T2 siRNA (m) is recommended for the inhibition of β-1,4-Gal-T2 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μM in 66 μl. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor β-1,4-Gal-T2 gene expression knockdown using RT-PCR Primer: β-1,4-Gal-T2 (m)-PR: sc-108224-PR (20 μl). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.