



RTRAF siRNA (m): sc-108820

BACKGROUND

RTRAF, also known as CGI-99, CLE, CLE7, C14orf166 or RLLM1, is a 244 amino acid protein that localizes to the nucleus, as well as to the cytoplasm, and belongs to the UPF0568 family. Expressed in a variety of tissues with particularly high levels in heart and skeletal muscle, RTRAF functions as a homodimer that interacts with Ninein and is thought to regulate Ninein function. RTRAF is overexpressed in brain tumor tissue, suggesting a role in tumor development and metastasis. The gene encoding RTRAF maps to human chromosome 14, which houses over 700 genes and comprises nearly 3.5% of the human genome. Chromosome 14 encodes the presenilin 1 (PSEN1) gene, which is one of the three key genes associated with the development of Alzheimer's disease (AD). The SERPINA1 gene is also located on chromosome 14 and, when defective, leads to the genetic disorder α 1-antitrypsin deficiency, which is characterized by severe lung complications and liver dysfunction.

REFERENCES

- Huarte, M., Sanz-Ezquerro, J.J., Roncal, F., Ortín, J. and Nieto, A. 2001. PA subunit from influenza virus polymerase complex interacts with a cellular protein with homology to a family of transcriptional activators. *J. Virol.* 75: 8597-8604.
- Howng, S.L., Hsu, H.C., Cheng, T.S., Lee, Y.L., Chang, L.K., Lu, P.J. and Hong, Y.R. 2004. A novel ninein-interaction protein, CGI-99, blocks ninein phosphorylation by GSK3 β and is highly expressed in brain tumors. *FEBS Lett.* 566: 162-168.
- Perez-González, A., Rodriguez, A., Huarte, M., Salanueva, I.J. and Nieto, A. 2006. hCLE/CGI-99, a human protein that interacts with the influenza virus polymerase, is a mRNA transcription modulator. *J. Mol. Biol.* 362: 887-900.
- Lupi, I., Broman, K.W., Tzou, S.C., Gutenberg, A., Martino, E. and Caturegli, P. 2008. Novel autoantigens in autoimmune hypophysitis. *Clin. Endocrinol.* 69: 269-278.
- Cui, Y., Wu, J., Zong, M., Song, G., Jia, Q., Jiang, J. and Han, J. 2009. Proteomic profiling in pancreatic cancer with and without lymph node metastasis. *Int. J. Cancer* 124: 1614-1621.
- Online Mendelian Inheritance in Man, OMIM™. 2009. Johns Hopkins University, Baltimore, MD. MIM Number: 610858. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: 2700060E02Rik (mouse) mapping to 14 A3.

PRODUCT

RTRAF siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see RTRAF shRNA Plasmid (m): sc-108820-SH and RTRAF shRNA (m) Lentiviral Particles: sc-108820-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

RTRAF siRNA (m) is recommended for the inhibition of RTRAF expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor RTRAF gene expression knockdown using RT-PCR Primer: RTRAF (m)-PR: sc-108820-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.