

ATXN7L3B siRNA (m): sc-108984

BACKGROUND

ATXN7L3B (putative ataxin-7-like protein 3B) is a 97 amino acid protein that belongs to the SGF11 family. The ATXN7L3B gene is conserved in chimpanzee, bovine and mouse, and maps to human chromosome 12q21.1. Encoding over 1,100 genes within 132 million bases, chromosome 12 makes up about 4.5% of the human genome. A number of skeletal deformities are linked to chromosome 12 including hypochondrogenesis, achondrogenesis and Kniest dysplasia. Noonan syndrome, which includes heart and facial developmental defects among the primary symptoms, is caused by a mutant form of PTPN11 gene product, SH-PTP2. Chromosome 12 is also home to a homeobox gene cluster, which encodes crucial transcription factors for morphogenesis, and the natural killer complex gene cluster encoding C-type lectin proteins that mediate the NK cell response to MHC I interaction.

REFERENCES

1. Strausberg, R.L., Feingold, E.A., Grouse, L.H., Derge, J.G., Klausner, R.D., Collins, F.S., Wagner, L., Shenmen, C.M., Schuler, G.D., Altschul, S.F., Zeeberg, B., Buetow, K.H., Schaefer, C.F., Bhat, N.K., Hopkins, R.F., et al. 2002. Generation and initial analysis of more than 15,000 full-length human and mouse cDNA sequences. *Proc. Natl. Acad. Sci. USA* 99: 16899-16903.
2. Ota, T., Suzuki, Y., Nishikawa, T., Otsuki, T., Sugiyama, T., Irie, R., Wakamatsu, A., Hayashi, K., Sato, H., Nagai, K., Kimura, K., Makita, H., Sekine, M., Obayashi, M., Nishi, T., Shibahara, T., Tanaka, T., et al. 2004. Complete sequencing and characterization of 21,243 full-length human cDNAs. *Nat. Genet.* 36: 40-45.
3. Walley, A.J., Asher, J.E. and Froguel, P. 2009. The genetic contribution to non-syndromic human obesity. *Nat. Rev. Genet.* 10: 431-442.
4. Guardiola, M.T., Dobin, S.M., Dal Cin, P. and Donner, L.R. 2010. Pericentric inversion (12)(p12q13-14) as the sole chromosomal abnormality in a leiomyoma of the vulva. *Cancer Genet. Cytogenet.* 199: 21-23.
5. Aytekin, T., Ozaslan, M. and Cengiz, B. 2010. Deletion mapping of chromosome region 12q13-24 in colorectal cancer. *Cancer Genet. Cytogenet.* 201: 32-38.
6. Ierardo, G., Luzzi, V., Panetta, F., Sfasciotti, G.L. and Polimeni, A. 2010. Noonan syndrome: a case report. *Eur. J. Paediatr. Dent.* 11: 97-100.
7. SWISS-PROT/TrEMBL (Q96GX2). World Wide Web URL: <http://www.uniprot.org/uniprot/Q96GX2>

CHROMOSOMAL LOCATION

Genetic locus: Atxn7l3b (mouse) mapping to 10 D2.

PRODUCT

ATXN7L3B siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ATXN7L3B shRNA Plasmid (m): sc-108984-SH and ATXN7L3B shRNA (m) Lentiviral Particles: sc-108984-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

ATXN7L3B siRNA (m) is recommended for the inhibition of ATXN7L3B expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ATXN7L3B gene expression knockdown using RT-PCR Primer: ATXN7L3B (m)-PR: sc-108984-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.