



# MAD2L1BP (m): 293T Lysate: sc-110183

## BACKGROUND

MAD2L1BP (MAD2L1 binding protein), also known as CMT2, is a 274 amino acid protein that localizes to the nucleoplasm during early mitosis and to the spindle from metaphase through anaphase. Functioning as a component of the spindle checkpoint (which delays the onset of anaphase until kinetochore attachment is complete), MAD2L1BP is thought to coordinate cell cycle events in late mitosis, possibly binding to MAD2, thereby silencing the spindle checkpoint and allowing mitosis to proceed. MAD2L1BP is expressed as multiple alternative spliced isoforms that, upon DNA damage, may be phosphorylated by ATM or ATR. The gene encoding MAD2L1BP maps to human chromosome 6, which contains 170 million base pairs and comprises nearly 6% of the human genome.

## REFERENCES

1. Nagase, T., Miyajima, N., Tanaka, A., Sazuka, T., Seki, N., Sato, S., Tabata, S., Ishikawa, K., Kawarabayashi, Y. and Kotani, H. 1995. Prediction of the coding sequences of unidentified human genes. III. The coding sequences of 40 new genes (KIAA0081-KIAA0120) deduced by analysis of cDNA clones from human cell line KG-1. *DNA Res.* 2: 37-43.
2. Howell, B.J., Hoffman, D.B., Fang, G., Murray, A.W. and Salmon, E.D. 2000. Visualization of Mad2 dynamics at kinetochores, along spindle fibers, and at spindle poles in living cells. *J. Cell Biol.* 150: 1233-1250.
3. Habu, T., Kim, S.H., Weinstein, J. and Matsumoto, T. 2002. Identification of a MAD2-binding protein, CMT2, and its role in mitosis. *EMBO J.* 21: 6419-6428.
4. Xia, G., Luo, X., Habu, T., Rizo, J., Matsumoto, T. and Yu, H. 2004. Conformation-specific binding of p31<sup>comet</sup> antagonizes the function of Mad2 in the spindle checkpoint. *EMBO J.* 23: 3133-3143.
5. Yang, M., Li, B., Tomchick, D.R., Machius, M., Rizo, J., Yu, H. and Luo, X. 2007. p31<sup>comet</sup> blocks Mad2 activation through structural mimicry. *Cell* 131: 744-755.
6. Yun, M.Y., Kim, S.B., Park, S., Han, C.J., Han, Y.H., Yoon, S.H., Kim, S.H., Kim, C.M., Choi, D.W., Cho, M.H., Park, G.H. and Lee, K.H. 2007. Mutation analysis of p31<sup>comet</sup> gene, a negative regulator of Mad2, in human hepatocellular carcinoma. *Exp. Mol. Med.* 39: 508-513.
7. Peters, J.M. 2008. Checkpoint control: the journey continues. *Curr. Biol.* 18: R170-R172.

## CHROMOSOMAL LOCATION

Genetic locus: Mad2l1bp (mouse) mapping to 17 C.

## PRODUCT

MAD2L1BP (m): 293T Lysate represents a lysate of mouse MAD2L1BP transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

## STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

## APPLICATIONS

MAD2L1BP (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive MAD2L1BP antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.