

IEX-1 (m2): 293T Lysate: sc-110242

BACKGROUND

Tumors are frequently observed as resistant to apoptotic induction by FAS, tumor necrosis factor α (TNF- α) or irradiation. This anti-death activity may be attributed to immediate early-response genes that are regulated at the transcriptional level, including the protein IEX-1. IEX-1 (immediately early gene X-1), also known as IER3 (immediate early response 3), DIF-2 (differentiation-dependent gene 2 protein), immediate early protein GLY96 or PRG1 (PACAP-responsive gene 1 protein), is a 156 amino acid single-pass type II membrane protein that belongs to the IER3 family. IEX-1 was originally characterized as a gene induced by ultraviolet radiation and TNF- α , which protected human squamous carcinoma cells from apoptosis. Subsequent transfection studies have also shown that expression of IEX-1 in human keratinocytes and mouse fibroblasts results in more rapid proliferation of the cells as compared with controls. The promoter region of IEX-1 contains binding motifs for both NF κ B and p53, suggesting that these proteins may regulate its expression.

REFERENCES

1. Beyaert, R., et al. 1994. Molecular mechanisms of tumor necrosis factor-induced cytotoxicity. What we do understand and what we do not. *FEBS Lett.* 340: 9-16.
2. Kondratyev, A.D., et al. 1996. Identification and characterization of a radiation-inducible glycosylated human early-response gene. *Cancer Res.* 56: 1498-1502.
3. Wu, M.X., et al. 1998. IEX-1, an apoptosis inhibitor involved in NF κ B-mediated cell survival. *Science* 281: 998-1001.
4. Kobayashi, T., et al. 1998. Regulation of a novel immediate early response gene, IEX-1, in keratinocytes by 1 α ,25-dihydroxyvitamin D3. *Biochem. Biophys. Res. Commun.* 251: 868-873.
5. Kumar, R., et al. 1998. A novel immediate early response gene, IEX-1, is induced by ultraviolet radiation in human keratinocytes. *Biochem. Biophys. Res. Commun.* 253: 336-341.
6. Schafer, H., et al. 1998. The promoter of human p22/PACAP response gene 1 (PRG1) contains functional binding sites for the p53 tumor suppressor and for NF κ B. *FEBS Lett.* 436: 139-143.

CHROMOSOMAL LOCATION

Genetic locus: Ier3 (mouse) mapping to 17 B1.

PRODUCT

IEX-1 (m2): 293T Lysate represents a lysate of mouse IEX-1 transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

IEX-1 (m2): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive IEX-1 antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.