



p41-ARCb (m): 293T Lysate: sc-110361

BACKGROUND

The Arp2/3 (Actin-related protein 2/3) complex consists of seven subunits, all of which are Actin-related proteins. The complex is involved in the control of Actin polymerization and in mediating the formation of branched Actin networks. The p41 subunit of Arp2/3 exists in multiple versions which arise due to posttranslational modifications and vary in function depending on cell type or developmental stage. p41-ARCb, also known as ARPC1B (Actin-related protein 2/3 complex subunit 1B) or ARC41 (Arp2/3 complex 41 kDa subunit), is a 372 amino acid version of the p41 subunit of the Arp2/3 complex. Localized to the cytoplasm and cytoskeleton, p41-ARCb is involved in regulating the assembly and maintenance of the complex and, unlike the other Arp2/3 subunits, is not primarily involved in Actin polymerization.

REFERENCES

1. Welch, M.D., DePace, A.H., Verma, S., Iwamatsu, A. and Mitchison, T.J. 1997. The human Arp2/3 complex is composed of evolutionarily conserved subunits and is localized to cellular regions of dynamic Actin filament assembly. *J. Cell Biol.* 138: 375-384.
2. Zhao, X., Yang, Z., Qian, M. and Zhu, X. 2001. Interactions among subunits of human Arp2/3 complex: p20-ARC as the hub. *Biochem. Biophys. Res. Commun.* 280: 513-517.
3. Kaneda, A., Kaminishi, M., Nakanishi, Y., Sugimura, T. and Ushijima, T. 2002. Reduced expression of the Insulin-induced protein 1 and p41 Arp2/3 complex genes in human gastric cancers. *Int. J. Cancer* 100: 57-62.
4. Vadlamudi, R.K., Li, F., Barnes, C.J., Bagheri-Yarmand, R. and Kumar, R. 2004. p41-ARC subunit of human Arp2/3 complex is a p21-activated kinase-1-interacting substrate. *EMBO Rep.* 5: 154-160.
5. Kaneda, A., Kaminishi, M., Sugimura, T. and Ushijima, T. 2004. Decreased expression of the seven Arp2/3 complex genes in human gastric cancers. *Cancer Lett.* 212: 203-210.
6. Kumagai, K., Nimura, Y., Mizota, A., Miyahara, N., Aoki, M., Furusawa, Y., Takiguchi, M., Yamamoto, S. and Seki, N. 2006. Arpc1b gene is a candidate prediction marker for choroidal malignant melanomas sensitive to radiotherapy. *Invest. Ophthalmol. Vis. Sci.* 47: 2300-2304.

CHROMOSOMAL LOCATION

Genetic locus: Arpc1b (mouse) mapping to 5 G2.

PRODUCT

p41-ARCb (m): 293T Lysate represents a lysate of mouse p41-ARCb transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

p41-ARCb (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive p41-ARCb antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.