

OCIAD1 (h): 293T Lysate: sc-110825

BACKGROUND

OCIAD1 (OCIA domain containing 1), also known as OCIA (ovarian carcinoma immunoreactive antigen), TPA018 or ASRIJ, is a 245 amino acid endosomal protein that contains one OCIA domain and belongs to the OCIAD1 family. OCIAD1 exists as two alternatively spliced isoforms (termed OCIAD1 isoform A and B) that localize to different tissues; isoform A localizes to prostate, brain, mammary gland, testis, placenta and ovary, whereas isoform B is restricted to the central nervous system. The gene encoding OCIAD1 maps to human chromosome 4, which represents approximately 6% of the human genome, contains nearly 900 genes and is associated with Huntington's disease, Ellis-van Creveld syndrome, methylmalonic acidemia and polycystic kidney disease.

REFERENCES

1. Kalchman, M.A., et al. 1996. Huntingtin is ubiquitinated and interacts with a specific ubiquitin-conjugating enzyme. *J. Biol. Chem.* 271: 19385-19394.
2. Howard, T.D., et al. 1997. Autosomal dominant postaxial polydactyly, nail dystrophy, and dental abnormalities map to chromosome 4p16, in the region containing the Ellis-van Creveld syndrome locus. *Am. J. Hum. Genet.* 61: 1405-1412.
3. Singhrao, S.K., et al. 1998. Huntingtin protein colocalizes with lesions of neurodegenerative diseases: an investigation in Huntington's, Alzheimer's, and Pick's diseases. *Exp. Neurol.* 150: 213-222.
4. Krakow, D., et al. 2000. Exclusion of the Ellis-van Creveld region on chromosome 4p16 in some families with asphyxiating thoracic dystrophy and short-rib polydactyly syndromes. *Eur. J. Hum. Genet.* 8: 645-648.
5. Sommardahl, C., et al. 2001. Phenotypic variations of orpk mutation and chromosomal localization of modifiers influencing kidney phenotype. *Physiol. Genomics* 7: 127-134.
6. Dobson, C.M., et al. 2002. Identification of the gene responsible for the cblA complementation group of vitamin B12-responsive methylmalonic acidemia based on analysis of prokaryotic gene arrangements. *Proc. Natl. Acad. Sci. USA* 99: 15554-15559.
7. Sengupta, S., et al. 2008. Ovarian cancer immuno-reactive antigen domain containing 1 (OCIAD1), a key player in ovarian cancer cell adhesion. *Gynecol. Oncol.* 109: 226-233.

CHROMOSOMAL LOCATION

Genetic locus: OCIAD1 (human) mapping to 4p11.

PRODUCT

OCIAD1 (h): 293T Lysate represents a lysate of human OCIAD1 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

OCIAD1 (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive OCIAD1 antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

OCIAD1 (E-12): sc-390906 is recommended as a positive control antibody for Western Blot analysis of enhanced human OCIAD1 expression in OCIAD1 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



OCIAD1 (E-12): sc-390906. Western blot analysis of OCIAD1 expression in non-transfected: sc-117752 (A) and human OCIAD1 transfected: sc-110825 (B) whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.