

Rpp29 (h): 293T Lysate: sc-111698

BACKGROUND

Ribonuclease P (PNase P) and Ribonuclease MRP (RNase MRP) are small nuclear ribonucleoproteins (snRNPs) that act on RNA substrates *in vitro*. RNase P and RNase MRP, which accumulate in the nucleolus, have a similar RNA component and also have several protein subunits in common. RNase P, which consists of a complex of an RNA species (H1 RNA), POP1 (processing of precursor 1), POP5 (processing of precursor 5), and at least 7 Rpps (including Rpp14, Rpp29, Rpp30 and Rpp38), removes the 5' leader sequences from precursor tRNA molecules. In particular, the nucleolar-localizing RNase P catalyzes the hydrolysis of a specific phosphodiester bond in precursor tRNA to form the mature 5' end of tRNA. The structurally related RNase MRP plays an essential role in the formation of the 5' end of 5.8S rRNA. Both RNase P and RNase MRP are associated with Th/To ribonucleoproteins; Rpp30 and Rpp38 have specifically been implicated as Th autoantigens which contribute to the autoimmune disease systemic sclerosis.

REFERENCES

1. Karwan, R. 1993. RNase MRP/RNase P: a structure-function relation conserved in evolution? FEBS Lett. 319: 1-4.
2. Jarrous, N., Eder, P.S., Guerrier-Takada, C., Hoog, C. and Altman, S. 1998. Autoantigenic properties of some protein subunits of catalytically active complexes of human ribonuclease P. RNA 4: 407-417.
3. Pluk, H., van Eenennaam, H., Rutjes, S.A., Pruijn, G.J. and van Venrooij, W.J. 1999. RNA-protein interactions in the human RNase MRP ribonucleoprotein complex. RNA 5: 512-524.
5. van Eenennaam, H., Jarrous, N., van Venrooij, W.J. and Pruijn, G.J. 2000. Architecture and function of the human endonucleases RNase P and RNase MRP. IUBMB Life 49: 265-272.
6. Altman, S. 2000. The road to RNase P. Nat. Struct. Biol. 7: 827-828.
7. Kurz, J.C. and Fierke, C.A. 2000. Ribonuclease P: a ribonucleoprotein enzyme. Curr. Opin. Chem. Biol. 4: 553-558.
8. van Eenennaam, H., van der Heijden, A., Janssen, R.J., van Venrooij, W.J. and Pruijn, G.J. 2001. Basic domains target protein subunits of the RNase MRP complex to the nucleolus independently of complex association. Mol. Biol. Cell 12: 3680-3689.

CHROMOSOMAL LOCATION

Genetic locus: POP4 (human) mapping to 19q12.

PRODUCT

Rpp29 (h): 293T Lysate represents a lysate of human Rpp29 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

Rpp29 (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive Rpp29 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.