

PGK1 (h2): 293T Lysate: sc-112139

BACKGROUND

Phosphoglycerate kinases 1/2 (PGK1/2), (ATP: 3-phospho-D-glycerate 1-phosphotransferase, EC 2.7.2.3) are somatically expressed, glycolytic enzymes that catalyze the transfer of a phosphoryl group from the acyl phosphate of 1,3-bisphosphoglycerate to ADP, thereby forming ATP and 3-phosphoglycerate. The human PGK gene is interrupted by 10 introns and spans 23 kilobases, and is X chromosome-linked at position Xq11-Xq13, a region implicated in prostate cancer, androgen insensitivity, perineal hypospadias, and other genetic abnormalities. In addition to influencing glycolysis, the PGK1 is secreted by tumor cells and contributes to proliferative angiogenic processes as a disulfide reductase. PGK1 mediated reduction of disulphide bonds in the serine proteinase plasmin initiates the release of the tumor blood vessel inhibitor angiostatin, an event that is critical for blood vessel formation or angiogenesis in tumor expansion and metastasis.

REFERENCES

1. Michelson, A.M., et al. 1985. Structure of the human phosphoglycerate kinase gene and the intron-mediated evolution and dispersal of the nucleotide-binding domain. *Proc. Natl. Acad. Sci. USA* 82: 6965-6969.
2. Ogino, T., et al. 1999. Involvement of a cellular glycolytic enzyme, phosphoglycerate kinase, in Sendai virus transcription. *J. Biol. Chem.* 274: 35999-36008.
3. Riley, D.E., et al. 1999. A hemizygous short tandem repeat polymorphism 3' to the human phosphoglycerate kinase gene. *Mol. Biol. Rep.* 26: 159-165.
4. Lay, A.J., et al. 2000. Phosphoglycerate kinase acts in tumour angiogenesis as a disulphide reductase. *Nature* 408: 869-873.
5. LocusLink Report (LocusID: 5230). <http://www.ncbi.nlm.nih.gov/LocusLink/>

CHROMOSOMAL LOCATION

Genetic locus: PGK1 (human) mapping to Xq21.1.

PRODUCT

PGK1 (h2): 293T Lysate represents a lysate of human PGK1 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

PGK1 (h2): 293T Lysate is suitable as a Western Blotting positive control for human reactive PGK1 antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

PGK1/2 (A-2): sc-166432 is recommended as a positive control antibody for Western Blot analysis of enhanced human PGK1 expression in PGK1 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



PGK1/2 (A-2): sc-166432. Western blot analysis of PGK1 expression in non-transfected: sc-117752 (A) and human PGK1 transfected: sc-112139 (B) 293T whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.