

# Thrombospondin 2 (h2): 293T Lysate: sc-112804

## BACKGROUND

The Thrombospondin proteins (TSP 1-4) compose a family of glycoproteins that are involved in cell-to-cell and cell-to-matrix signaling. These extracellular, cell-surface proteins form complexes of both homo- and heteromultimers. Thrombospondins play a role in development, aggregation of platelets, adhesion and migration of cells, and progression of cells through the growth cycle. Thrombospondin 1 is released from platelets in response to Thrombin stimulation and is a transient component of the extracellular matrix of developing and repairing tissues. Thrombospondin 2 shares a high degree of homology with Thrombospondin 1 and is thought to have overlapping but unique functions. Thrombospondin 3 is a developmentally regulated heparin binding protein. Thrombospondin 4 is neuronally expressed and stimulates neurite outgrowth.

## REFERENCES

1. Mosher, D.F. 1990. Physiology of Thrombospondin. *Annu. Rev. Med.* 41: 85-97.
2. Bornstein, P., O'Rourke, K., Wilkstrom, K., Wolf, F.W., Katz, R., Li, P. and Dixit, V.M. 1991. A second, expressed Thrombospondin gene (Thbs2) exists in the mouse genome. *J. Biol. Chem.* 266: 12821-12824.
3. LaBell, T.L., Milewicz, D.J., Distech, C.M. and Byers, B.H. 1992. Thrombospondin II: partial cDNA sequence, chromosome location and expression of a second member of the Thrombospondin gene family in humans. *Genomics* 12: 421-429.
4. O'Rourke, K.M., Laherty, C.D. and Dixit, V.M. 1992. Thrombospondin 1 and Thrombospondin 2 are expressed as both homo- and heterotrimers. *J. Biol. Chem.* 267: 24921-24924.
5. Jahav, J. 1993. The functions of Thrombospondin and its involvement in physiology and pathophysiology. *Biochem. Biophys. Acta* 1182: 1-14.
6. Arber, S. and Caroni, P. 1995. Thrombospondin 4, an extracellular matrix protein expressed in the developing and adult nervous system promotes neurite outgrowth. *J. Cell Biol.* 131: 1083-1094.
7. Qabar, A., Derick, L., Lawler, J. and Dixit, V. 1995. Thrombospondin 3 is a pentameric molecule held together by interchain disulfide linkage involving two cysteine residues. *J. Biol. Chem.* 270: 12725-12729.
8. Adams, J.C. 1997. Thrombospondin 1. *Int. J. Biochem. Cell Biol.* 29: 861-865.

## CHROMOSOMAL LOCATION

Genetic locus: THBS2 (human) mapping to 6q27.

## PRODUCT

Thrombospondin 2 (h2): 293T Lysate represents a lysate of human Thrombospondin 2 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

## STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

## APPLICATIONS

Thrombospondin 2 (h2): 293T Lysate is suitable as a Western Blotting positive control for human reactive Thrombospondin 2 antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

Thrombospondin 1/2 (D-9): sc-74538 is recommended as a positive control antibody for Western Blot analysis of enhanced human Thrombospondin 2 expression in Thrombospondin 2 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:  
1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

## DATA



Thrombospondin 1/2 (D-9): sc-74538. Western blot analysis of Thrombospondin 2 expression in non-transfected: sc-117752 (A) and truncated human Thrombospondin 2 transfected: sc-112804 (B) 293T whole cell lysates.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.