

CD32 (h2): 293T Lysate: sc-112840

BACKGROUND

CD32 (also designated Fc γ RII) is a low affinity receptor for the Fc fragment of aggregated IgG. CD32 is responsible for the clearance of immunocomplexes by macrophages and also plays an important role in the regulation of antibody production by B cells. IgG can noncooperatively bind either one or two highly glycosylated CD32 molecules and this binding delivers a negative signal for B cells. CD32 exists as several isoforms that are produced by alternative splicing of three distinct genes, A, B and C. These isoforms are designated Fc γ RIIA, Fc γ RIIB1, Fc γ RIIB3 and Fc γ RIIC. All isoforms are present on monocytes, placental trophoblasts and endothelial cells. In addition, the Fc γ RIIB forms are present on B lymphocytes, and the Fc γ RIIA and Fc γ RIIC forms are found on neutrophils.

REFERENCES

1. Bijsterbosch, M.K. and Klaus, G.G. 1985. Crosslinking of surface immunoglobulin and Fc receptors on B lymphocytes inhibits stimulation of inositol phospholipid breakdown via the antigen receptors. *J. Exp. Med.* 162: 1825-1836.
2. Huizinga, T.W.J., Kerst, M., Nuyens, J.H., Vlug, A., von dem Borne, A.E., Roos, D. and Tetteroo, P.A. 1989. Binding characteristics of dimeric IgG subclass complexes to human neutrophils. *J. Immunol.* 142: 2365-2369.
3. Stuart, S.G., Simister, N.E., Clarkson, S.B., Kacinski, B.M., Shapiro, M. and Mellman, I. 1989. Human IgG Fc receptor (hFcRII; CD32) exists as multiple isoforms in macrophages, lymphocytes and IgG-transporting placental epithelium. *EMBO J.* 8: 3657-3666.
4. Raveth, J.V. and Kinet, J.P. 1991. Fc Receptors. *Annu. Rev. Immunol.* 9: 457-492.
5. Barclay, A.N., Beyers, A.D., Birkeland, M.L., Brown, S.J., Somoza, C. and Williams, A.F. 1993. *The leukocyte antigen facts book*. London: Academic Press, 170-172.
6. Sondermann, P., Jacob, U., Kutscher, C. and Frey, J. 1999. Characterization and crystallization of soluble human Fc γ receptor II (CD32) isoforms produced in insect cells. *Biochemistry* 38: 8469-8477.

CHROMOSOMAL LOCATION

Genetic locus: FCGR2B (human) mapping to 1q23.3.

PRODUCT

CD32 (h2): 293T Lysate represents a lysate of human CD32 transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

APPLICATIONS

CD32 (h2): 293T Lysate is suitable as a Western Blotting positive control for human reactive CD32 antibodies.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.