

CD40 (h): 293 Lysate: sc-112948

BACKGROUND

Resting B cells can be activated and clonally expanded into antibody-producing cells in response to a combination of cell contact and soluble signals provided by primed helper T (Th) cells. While cytokines IL-4 and IL-13 alone are inadequate for B cell activation, contact with Th cells seems to be sufficient for delivery of proliferative signals. A receptor ligand pair central to the transmission of this signal is CD40, expressed on the surface of B cells, together with CD40L, expressed on activated T cells. In the presence of such stimulus, IL-4 and IL-13 are capable of triggering immunoglobulin class switching and secretion of IgE. B cells are sensitive to these cytokines only subsequent to CD40/CD40L-driven DNA synthesis. A downstream mediator of the CD40 signaling pathway, designated CRAF, is a member of an expanding family of proteins that contain a conserved cysteine- and histidine-rich RING finger motif. Other members of the family include TRAF1 and TRAF2. The latter proteins have been shown to regulate TNF-R2 as well as CD40 signaling through activation of the NF κ B family of transcription factors.

REFERENCES

1. Kehry, M.R. and Hodgkin, P.D. 1994. B cell activation by helper T cell membranes. *Crit. Rev. Immunol.* 14: 221-238.
2. Hu, H.M., O'Rourke, K., Boguski, M.S. and Dixit, V.M. 1994. A novel RING finger protein interacts with the cytoplasmic domain of CD40. *J. Biol. Chem.* 269: 30069-30072.
3. Rothe, M., Wong, S.C., Henzel, W.J. and Goeddel, D.V. 1994. A novel family of putative signal transducers associated with the cytoplasmic domain of the 75 kDa tumor necrosis factor receptor. *Cell* 78: 681-682.
4. Gordon, J. 1995. CD40 and its ligand: central players in B lymphocyte survival, growth and differentiation. *Blood Rev.* 9: 53-56.
5. Fuleihan, R., Ahern, D. and Geha, R.S. 1995. Expression of the CD40 ligand in T lymphocytes and induction of IgE isotype switching. *Int. Arch. Allergy Immunol.* 107: 43-44.
6. Cheng, G., Cleary, A.M., Ye, Z.S., Hong, D.I., Lederman, S. and Baltimore, D. 1995. Involvement of CRAF1, a relative of TRAF, in CD40 signaling. *Science* 267: 1494-1498.
7. Rothe, M., Sarma, V., Dixit, V.M. and Goeddel, D.V. 1995. TRAF2-mediated activation of NF κ B by TNF receptor 2 and CD40. *Science* 269: 1424-1427.
8. Grandien, A., Brás, A. and Martinez, C. 1996. Acquisition of CD40 expression during murine B cell differentiation. *Scand. J. Immunol.* 43: 47-55.

CHROMOSOMAL LOCATION

Genetic locus: CD40 (human) mapping to 20q13.12.

PRODUCT

CD40 (h): 293 Lysate represents a lysate of human CD40 transfected 293 cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

CD40 (h): 293 Lysate is suitable as a Western Blotting positive control for human reactive CD40 antibodies. Recommended use: 10-20 μ l per lane.


Control 293 Lysate: sc-110760 is available as a Western Blotting negative control lysate derived from non-transfected 293 cells.

CD40 (H-10): sc-13128 is recommended as a positive control antibody for Western Blot analysis of enhanced human CD40 expression in CD40 transfected 293 cells (starting dilution 1:100, dilution range 1:100-1:1,000).


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048.

DATA



CD40 (H-10): sc-13128. Near-infrared western blot analysis of CD40 expression in non-transfected: sc-110760 (A) and human CD40 transfected: sc-112948 (B) whole cell lysates. Blocked with UltraCruz® Blocking Reagent: sc-516214. Detection reagent used: m-IgG κ BP-CFL 680: sc-516180.



CD40 (LOB-11): sc-13528. Western blot analysis of CD40 expression in non-transfected: sc-110760 (A) and human CD40 transfected: sc-112948 (B) 293 whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.