



CD9 (h3): 293T Lysate: sc-113009

BACKGROUND

CD9 is a type IV transmembrane glycoprotein with four transmembrane domains. CD9 on pre-B cells may play a role in cell-cell adhesion. In addition, CD9 may play a role in signal transduction mediated by interaction with low molecular weight GTP binding proteins. CD9 is expressed on early B cells, eosinophils, basophils and activated T cells and is a major component of the platelet cell surface. It is also expressed on most non-T acute lymphoblastic leukemia cells and on some acute myeloid and chronic lymphoid leukemias.

REFERENCES

1. Seehafer, J.G., et al. 1991. Evidence that the signal-initiating membrane protein CD9 is associated with small GTP-binding proteins. *Biochem. Biophys. Res. Commun.* 179: 401-406.
2. Lanza, F., et al. 1991. cDNA cloning and expression of platelet p24/CD9. Evidence for a new family of multiple membrane-spanning proteins. *J. Biol. Chem.* 266: 10638-10645.
3. Ferrero, D., et al. 1991. CD9 antigen on acute non-lymphoid leukemia cells: preferential expression by promyelocytic (M3) subtype. *Leuk. Res.* 15: 457-461.
4. Wright, M.D., et al. 1994. The ins and outs of the transmembrane 4 superfamily. *Immunol. Today* 15: 588-594.
5. Masellis-Smith, A., et al. 1994. CD9-regulated adhesion. Anti-CD9 monoclonal antibody induce pre-B cell adhesion to bone marrow fibroblasts through *de novo* recognition of Fibronectin. *J. Immunol.* 152: 2768-2777.
6. Xu, M., et al. 1994. Regulation of CD9 expression during 12-O-tetradecanoyl-phorbol-13-acetate-induced differentiation of human Myeloid leukemia (HL-60) cells. *Cell Growth Differ.* 5: 1225-1234.
7. Nakamura, K., et al. 1995. Membrane-anchored heparin-binding EGF-like growth factor (HB-EGF) and diphtheria toxin receptor-associated protein (DRAP27)/CD9 form a complex with Integrin $\alpha 3\beta 1$ at cell-cell contact sites. *J. Cell Biol.* 129: 1691-1705.
8. Slupsky, J.R., et al. 1997. Analysis of CD9, CD32 and p67 signalling; use of degranulated platelets indicates direct involvement of CD9 and p67 in integrin activation. *Br. J. Haematol.* 96: 275-286.
9. Le Naour, F., et al. 1997. Upregulation of CD9 expression during TPA treatment of K-562 cells. *Leukemia* 11: 1290-1297.

CHROMOSOMAL LOCATION

Genetic locus: CD9 (human) mapping to 12p13.31.

PRODUCT

CD9 (h3): 293T Lysate represents a lysate of human CD9 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

CD9 (h3): 293T Lysate is suitable as a Western Blotting positive control for human reactive CD9 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.