

SRPK2 (h): 293T Lysate: sc-113878

BACKGROUND

Arginine/serine-rich (RS) domain-containing proteins and their phosphorylation by specific protein kinases constitute control circuits to regulate both constitutive and alternative pre-mRNA splicing and coordinate splicing with transcription in cells. Two SR protein-specific kinases (SRPK, also designated SFRSK), SRPK1 and SRPK2, are highly specific for the phosphorylation of these RS proteins, thereby contributing to splicing regulation. SRPK1 plays a role in the condensation of sperm chromatin. SRPK2 has a stringent preference for SR dipeptides and contains a proline-rich sequence at its amino-terminus. SRPK1 is expressed as two alternatively spliced isoforms, one of which is localized specifically to testis and the other of which shares a similar localization pattern to SRPK2 and is present in brain, heart, liver and lung.

REFERENCES

1. Wang, H.Y., et al. 1998. SRPK2: a differentially expressed SR protein-specific kinase involved in mediating the interaction and localization of pre-mRNA splicing factors in mammalian cells. *J. Cell Biol.* 140: 737-750.
2. Kuroyanagi, N., et al. 1998. Novel SR-protein-specific kinase, SRPK2, disassembles nuclear speckles. *Biochem. Biophys. Res. Commun.* 242: 357-364.
3. Papoutsopoulou, S., et al. 1999. SR protein-specific kinase 1 is highly expressed in testis and phosphorylates protamine 1. *Nucleic Acids Res.* 27: 2972-2980.
4. Wang, H.Y., et al. 1999. Localization of serine kinases, SRPK1 (SFRSK1) and SRPK2 (SFRSK2), specific for the SR family of splicing factors in mouse and human chromosomes. *Genomics* 57: 310-315.
5. Tang, Z., et al. 2000. Biochemical and genetic conservation of fission yeast Dsk1 and human SR protein-specific kinase 1. *Mol. Cell. Biol.* 20: 816-824.

CHROMOSOMAL LOCATION

Genetic locus: SRPK2 (human) mapping to 7q22.3.

PRODUCT

SRPK2 (h): 293T Lysate represents a lysate of human SRPK2 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

SRPK2 (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive SRPK2 antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

SRPK2 (A-5): sc-390930 is recommended as a positive control antibody for Western Blot analysis of enhanced human SRPK2 expression in SRPK2 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



SRPK2 (A-5): sc-390930. Western blot analysis of SRPK2 expression in non-transfected: sc-117752 (A) and human SRPK2 transfected: sc-113878 (B) 293T whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.