

15-LO (h): 293T Lysate: sc-114262

BACKGROUND

Lipoxygenases (LOs) are a family of enzymes which dioxygenate unsaturated fatty acids, thus initiating lipoperoxidation of membranes, the synthesis of signalling molecules, as well as inducing structural and metabolic changes in the cell. The Lox enzymes in mammals, 12-LO and 15-LO, are classified with respect to their positional specificity of the deoxygenation of their most common substrate, arachidonic acid. The metabolism of arachidonic acid leads to the generation of biologically active metabolites that have been implicated in cell growth and proliferation, as well as survival and apoptosis. 15-LO acts in physiological membrane remodeling and the pathogenesis of atherosclerosis, inflammation and carcinogenesis. It is highly regulated and expressed in a tissue- and cell-type-specific fashion. IL-4 and IL-13 play important roles in transactivating the 15-LO gene. Overexpression of 15-LO type 1 in prostate cancer contributes to the cancer progression by regulating IGF-1R expression and activation.

REFERENCES

1. Fletcher-Cieutat, M., Vanderhoek, J.Y., Bryant, R.W. and Bailey, J.M. 1985. Aspirin enhances the sensitivity of human platelet 12-lipoxygenase to inhibition by 15-HETE, an endogenous regulator. *Prostaglandins Leukot. Med.* 18: 255-259.
2. Pidgeon, G.P., Tang, K., Cai, Y.L., Piasentin, E. and Honn, K.V. 2003. Overexpression of platelet-type 12-lipoxygenase promotes tumor cell survival by enhancing $\alpha V/\beta 3$ and $\alpha V/\beta 5$ Integrin expression. *Cancer Res.* 63: 4258-4267.
3. Liu, C., Xu, D., Sjöberg, J., Forsell, P., Björkholm, M. and Claesson, H.E. 2004. Transcriptional regulation of 15-lipoxygenase expression by promoter methylation. *Exp. Cell Res.* 297: 61-67.
4. Kelavkar, U.P. and Cohen, C. 2004. 15-lipoxygenase-1 expression upregulates and activates Insulin-like growth factor-1 receptor in prostate cancer cells. *Neoplasia* 6: 41-52.
5. Raso, E., Döme, B., Somlai, B., Zacharek, A., Haggmann, W., Honn, K.V. and Tímár, J. 2004. Molecular identification, localization and function of platelet-type 12-lipoxygenase in human melanoma progression, under experimental and clinical conditions. *Melanoma Res.* 14: 245-250.

CHROMOSOMAL LOCATION

Genetic locus: ALOX15 (human) mapping to 17p13.2.

PRODUCT

15-LO (h): 293T Lysate represents a lysate of human 15-LO transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

APPLICATIONS

15-LO (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive 15-LO antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.