

P-Selectin (h): 293T Lysate: sc-114655

BACKGROUND

Selectins, also designated CD62 antigens, comprise a family of carbohydrate-binding proteins involved in mediating cellular interactions with leukocytes. L-Selectin (also designated LECAM-1 or CD62L) is expressed on the majority of B and naive T cells and on most monocytes, neutrophils and eosinophils. L-Selectin interacts with specific carbohydrates expressed by activated endothelial cells. P-Selectin (also designated GMP-140 or CD62P), expressed on activated platelets and endothelial cells, and E-Selectin (also designated ELMA-1 or CD62E), expressed on endothelial cells, exhibit overlapping ligand specificities. Both recognize sialyl-Le^x as a ligand and bind to specific carbohydrates on neutrophils and monocytes.

REFERENCES

1. Varki, A. 1994. Selectin ligands. *Proc. Natl. Acad. Sci. USA* 91: 7390-7397.
2. Tedder, T.F., Steeber, D.A., Chen, A. and Engel, P. 1995. The selectins: vascular adhesion molecules. *FASEB J.* 10: 866-873.
3. Lasky, L.A. 1995. Selectin-carbohydrate interactions and the initiation of the inflammatory response. *Annu. Rev. Biochem.* 64: 113-139.
4. Pavalko, F.M., Walker, D.M., Graham, L., Goheen, M., Doerschuk, C.M. and Kansas, G.S. 1995. The cytoplasmic domain of L-Selectin interacts with cytoskeletal proteins via α -Actinin: receptor positioning in microvilli does not require interaction with α -Actinin. *J. Cell Biol.* 129: 1155-1164.
5. Rosen, S.D. and Bertozzi, C.R. 1996. Two selectins converge on sulphate. *Leukocyte adhesion. Curr. Biol.* 6: 261-264.
6. McEver, R.P., Moore, K.L. and Cummings, R.D. 1996. Leukocyte trafficking mediated by selectin-carbohydrate interactions. *J. Biol. Chem.* 270: 11025-11028.
7. Diacovo, T.G., Puri, K.D., Warnock, R.A., Springer, T.A. and von Andrian, U.H. 1996. Platelet-mediated lymphocyte delivery to high endothelial venules. *Science* 273: 252-255.
8. Frenette, P.S., Mayadas, T.N., Rayburn, H., Hynes, R.O. and Wagner, D.D. 1996. Susceptibility to infection and altered hematopoiesis in mice deficient in both P- and E-Selectins. *Cell* 84: 563-574.

CHROMOSOMAL LOCATION

Genetic locus: SELP (human) mapping to 1q24.2.

PRODUCT

P-Selectin (h): 293T Lysate represents a lysate of human P-Selectin transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

P-Selectin (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive P-Selectin antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.