



Per1 (h): 293T Lysate: sc-114656

BACKGROUND

Biological timepieces called circadian clocks are responsible for the regulation of hormonal rhythms, sleep cycles and other behaviors. The suprachiasmatic nucleus (SCN), which is located in the brain, was the first mammalian circadian clock to be discovered. A number of transcription factors appearing to be molecular components of the SCN clock have been identified. Mutations within the Clock gene increase the length of the endogenous period and cause a loss of rhythmicity of circadian oscillations. Three mammalian period proteins, designated Per1, Per2 and Per3, exhibit circadian rhythms in the SCN. During subjective night, Per1 and Per2 RNA levels increase in response to light pulses while Per3 RNA levels show no change in response to light pulses. Tim, for timeless, interacts with Per1 as well as Per2; and Tim and Per1 negatively regulate Clock-BMAL1-induced transcription. Per1 protein isoforms display discrete cellular compartmentalization as well as tissue-specific size differences. The full size Per1 isoform is found principally in the cytoplasm while a shorter nuclear isoform also exists.

REFERENCES

1. Morell, V. 1995. A 24-hour circadian clock is found in the mammalian retina. *Science* 272: 349.
2. King, D.P., Vitaterna, M.H., Chang, A.M., Dove, W.F., Pinto, L.H., Turek, F.W. and Takahashi, J.S. 1997. The mouse Clock mutation behaves as an antimorph and maps within the W19H deletion, distal of Kit. *Genetics* 146: 1049-1060.
3. Antoch, M.P., Song, E.J., Chang, A.M., Vitaterna, M.H., Zha, Y., Wilsbacher, L.D., Sangoram, A.M., King, D.P., Pinto, L.H. and Takahashi, J.S. 1997. Functional identification of the mouse circadian Clock gene by transgenic BAC rescue. *Cell* 89: 655-667.
4. Zylka, M.J., Shearman, L.P., Weaver, D.R. and Reppert, S.M. 1998. Three period homologs in mammals: differential light responses in the suprachiasmatic circadian clock and oscillating transcripts outside of brain. *Neuron* 20: 1103-1110.
5. Sangoram, A.M., Saez, L., Antoch, M.P., Gekakis, N., Staknis, D., Whiteley, A., Fruechte, E.M., Vitaterna, M.H., Shimomura, K., King, D.P., Young, M.W., Weitz, C.J. and Takahashi, J.S. 1998. Mammalian circadian autoregulatory loop: a timeless ortholog and mPer1 interact and negatively regulate CLOCK-BMAL1-induced transcription. *Neuron* 21: 1101-1113.
6. Chilov, D., Hofer, T., Bauer, C., Wenger, R.H. and Gassmann, M. 2001. Hypoxia affects expression of circadian genes PER1 and CLOCK in mouse brain. *FASEB J.* 15: 2613-2622.

CHROMOSOMAL LOCATION

Genetic locus: PER1 (human) mapping to 17p13.1.

PRODUCT

Per1 (h): 293T Lysate represents a lysate of human Per1 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

Per1 (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive Per1 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.