



CD94 (h): 293T Lysate: sc-114690

BACKGROUND

The activity of natural killer (NK) cells is regulated by members of multiple receptor families that recognize class I MHC molecules, such as the killer cell inhibitory receptor/leukocyte immunoglobulin-like receptor (KIR/LIR) family and the C-type lectin superfamily. The KIR/LIR family includes p91A (also designated pp130 or PIR-B, for paired immunoglobulin-like receptor-B) and p91B (also designated PIR-A). p91A acts as an inhibitory receptor through interactions with SHP-1, whereas p91B acts as an activating receptor. CD94, NKG2 and Ly-49 are members of the C-type lectin superfamily of type II membrane glycoproteins. CD94 forms heterodimers with NKG2 isoforms on the surface of NK cells, whereas Ly-49 isoforms form homodimers. NKG2-D, expressed on NK cells, $\gamma\delta$ T cells, and CD8 $^+$ $\alpha\beta$ T cells, is a receptor for the stress inducible protein MICA, an antigen frequently expressed in epithelial tumors.

REFERENCES

- Long, E.O. and Wagtmann, N. 1997. Natural killer cell receptors. *Curr. Opin. Immunol.* 9: 344-350.
- Moretta, A. and Moretta, L. 1997. HLA class I specific inhibitory receptors. *Curr. Opin. Immunol.* 9: 694-701.
- Hayami, K., Fukuta, D., Nishikawa, Y., Yamashita, Y., Inui, M., Ohyama, Y., Hikida, M., Ohmori, H. and Takai, T. 1997. Molecular cloning of a novel murine cell-surface glycoprotein homologous to killer cell inhibitory receptors. *J. Biol. Chem.* 272: 7320-7327.
- Ryan, J.C. and Seaman, W.E. 1997. Divergent functions of lectin-like receptors on NK cells. *Immunol. Rev.* 155: 79-89.
- Vance, R.E., Tanamachi, D.M., Hanke, T. and Raulet, D.H. 1997. Cloning of a mouse homolog of CD94 extends the family of C-type lectins on murine natural killer cells. *Eur. J. Immunol.* 27: 3236-3241.
- Berg, K.L., Carlberg, K., Rohrschneider, L.R., Siminovitch, K.A. and Stanley, E.R. 1998. The major SHP-1-binding, tyrosine-phosphorylated protein in macrophages is a member of the KIR/LIR family and an SHP-1 substrate. *Oncogene* 17: 2535-2541.
- Salcedo, M. 1999. Inhibitory role of murine Ly-49 lectin-like receptors on natural killer cells. *Curr. Top. Microbiol. Immunol.* 244: 97-105.
- Bauer, S., Groh, V., Wu, J., Steinle, A., Phillips, J.H., Lanier, L.L. and Spies, T. 1999. Activation of NK cells and T cells by NKG2D, a receptor for stress-inducible MICA. *Science* 285: 727-729.
- Vance, R.E., Jamieson, A.M. and Raulet, D.H. 1999. Recognition of the class Ib molecule Qa-1 b by putative activating receptors CD94/NKG2C and CD94/NKG2E on mouse natural killer cells. *J. Exp. Med.* 190: 1801-1812.

CHROMOSOMAL LOCATION

Genetic locus: KLRD1 (human) mapping to 12p13.2.

PRODUCT

CD94 (h): 293T Lysate represents a lysate of human CD94 transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

CD94 (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive CD94 antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.