

NNT (h): 293T Lysate: sc-114805

BACKGROUND

The process of cellular respiration is carried out by integral inner mitochondrial membrane proteins that facilitate the harnessing of energy released by the oxidation of NADH. NNT (nicotinamide nucleotide transhydrogenase), also known as mitochondrial NAD(P) transhydrogenase or pyridine nucleotide *trans*-hydrogenase, is a 1,086 amino acid multi-pass mitochondrial inner membrane protein. NNT is a homodimer with an N-terminal section belonging to the AlaDH/PNT family and a C-terminal section belonging to the PNT β subunit family. NNT catalyzes the transfer of a hydride ion from NADH to NADP⁺ and functions as a mitochondrial inner membrane proton pump. Using the energy of the proton gradient created by the electron transport chain, NNT produces high concentrations of NADPH, which is used in free radical detoxification and biosynthesis.

REFERENCES

1. Forsmark-Andrée, P., Persson, B., Radi, R., Dallner, G. and Ernster, L. 1996. Oxidative modification of nicotinamide nucleotide transhydrogenase in sub-mitochondrial particles: effect of endogenous ubiquinol. *Arch. Biochem. Biophys.* 336: 113-120.
2. Arkblad, E.L., Betsholtz, C. and Rydström, J. 1996. The cDNA sequence of proton-pumping nicotinamide nucleotide transhydrogenase from man and mouse. *Biochim. Biophys. Acta* 1273: 203-205.
3. Zieger, B. and Ware, J. 1997. Cloning and deduced amino acid sequence of human nicotinamide nucleotide transhydrogenase. *DNA Seq.* 7: 369-373.
4. White, S.A., Peake, S.J., McSweeney, S., Leonard, G., Cotton, N.P. and Jackson, J.B. 2000. The high-resolution structure of the NAD(PH)-binding component (DIII) of proton-translocating transhydrogenase from human heart mitochondria. *Structure* 8: 1-12.
5. Arkblad, E.L., Egorov, M., Shakhparonov, M., Romanova, L., Polzikov, M. and Rydström, J. 2002. Expression of proton-pumping nicotinamide nucleotide transhydrogenase in mouse, human brain and *C. elegans*. *Comp. Biochem. Physiol. B, Biochem. Mol. Biol.* 133: 13-21.
6. Jackson, J.B. 2003. Proton translocation by transhydrogenase. *FEBS Lett.* 545: 18-24.
7. Online Mendelian Inheritance in Man, OMIM[™]. 2003. Johns Hopkins University, Baltimore, MD. MIM Number: 607878. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: NNT (human) mapping to 5p12.

PRODUCT

NNT (h): 293T Lysate represents a lysate of human NNT transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

NNT (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive NNT antibodies. Recommended use: 10-20 μ l per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

NNT (43-R17): sc-100490 is recommended as a positive control antibody for Western Blot analysis of enhanced human NNT expression in NNT transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



NNT (43-R17): sc-100490. Western blot analysis of NNT expression in non-transfected: sc-117752 (A) and human NNT transfected: sc-114805 (B) 293T whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.