

SERCA2 (h): 293T Lysate: sc-115268

BACKGROUND

ATP dependent calcium pumps are responsible, in part, for the maintenance of low cytoplasmic free calcium concentrations. The ATP pumps that reside in intracellular organelles are encoded by a family of structurally related enzymes, termed the sarcoplasmic or endoplasmic reticulum calcium (SERCA) ATPases. The sarcoplasmic reticulum of striated muscle is a specialized intracellular membrane system that plays a critical role in the contraction and relaxation of muscle. The SERCAs mediate Ca^{2+} uptake into intracellular stores. SERCA-mediated Ca^{2+} uptake induces and maintains muscular relaxation. The SERCA1 gene is exclusively expressed in type II (fast) skeletal muscle. The SERCA2 gene is subject to tissue-dependent processing which is responsible for the generation of the SERCA2a muscle-specific form expressed in type I (slow) skeletal, cardiac and smooth muscle and the SERCA2b isoform expressed in all cell types. The SERCA3 gene is not as well characterized and is found in non-muscle cells. SERCA2 plays an important part in regulating cardiac contractile function. SERCA3 is an isoform expressed in several cell types including platelets, lymphoid cells and mast cells. SERCA1, 2 and 3 all undergo alternative splicing.

REFERENCES

1. Aubier, M. and Viires, N. 1998. Calcium ATPase and respiratory muscle function. *Eur. Respir. J.* 11: 758-766.
2. Anger, M., Lompré, A.M., Vallot, O., Marotte, F., Rappaport, L. and Samuel, J.L. 1998. Cellular distribution of Ca^{2+} pumps and Ca^{2+} release channels in rat cardiac hypertrophy induced by aortic stenosis. *Circulation* 98: 2477-2486.
3. Loukianov, E., Ji, Y., Grupp, I.L., Kirkpatrick, D.L., Baker, D.L., Loukianova, T., Grupp, G., Lytton, J., Walsh, R.A. and Periasamy, M. 1998. Enhanced myo-cardial contractility and increased Ca^{2+} transport function in transgenic hearts expressing the fast-twitch skeletal muscle sarcoplasmic reticulum Ca^{2+} -ATPase. *Circ. Res.* 83: 889-897.
4. Bobe, R., Lacabartz-Porret, C., Bredoux, R., Martin, V., Ozog, A., Launay, S., Corvazier, E., Kovács, T., Papp, B. and Enouf, J. 1998. Expression of two isoforms of the third sarco/endoplasmic reticulum Ca^{2+} ATPase (SERCA3) in platelets. Possible recognition of the SERCA3b isoform by the PL/IM430 monoclonal antibody. *FEBS Lett.* 423: 259-264.
5. Poch, E., Leach, S., Snape, S., Cacic, T., MacLennan, D.H. and Lytton, J. 1998. Functional characterization of alternatively spliced human SERCA3 transcripts. *Am. J. Physiol., Cell Physiol.* 275: C1449-C1458.
6. Ozog, A., Pouzet, B., Bobe, R. and Lompré, A.M. 1998. Characterization of the 3' end of the mouse SERCA3 gene and tissue distribution of mRNA spliced variants. *FEBS Lett.* 427: 349-352.

CHROMOSOMAL LOCATION

Genetic locus: ATP2A2 (human) mapping to 12q24.11.

PRODUCT

SERCA2 (h): 293T Lysate represents a lysate of human SERCA2 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

SERCA2 (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive SERCA2 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.