

XTRP2 (h): 293T Lysate: sc-115985

BACKGROUND

The solute carrier (SLC) family, also known as the neurotransmitter transporter family, is one of the largest transporter families in the human genome. The SLC6 subgroup, which includes the dopamine transporter (DAT), the serotonin transporter (SERT) and the norepinephrine transporter (NET), contains sodium/chloride dependent, high-affinity plasma transport proteins. Sodium/chloride-dependent transporter XTRP2, also known as solute carrier family 6 member 18 (SLC6A18), is a 628 amino acid member of the SLC family. XTRP2 is composed of 12 exons, 11 introns and 8 tandem repeats. Highly expressed in the kidney, SLC6A18 is classified as an orphan transporter.

REFERENCES

1. Nash, S.R., Giros, B., Kingsmore, S.F., Kim, K.M., el-Mestikawy, S., Dong, Q., Fumagalli, F., Seldin, M.F. and Caron, M.G. 1998. Cloning, gene structure and genomic localization of an orphan transporter from mouse kidney with six alternatively-spliced isoforms. *Receptors Channels* 6: 113-128.
2. Quan, H., Athirakul, K., Wetsel, W.C., Torres, G.E., Stevens, R., Chen, Y.T., Coffman, T.M. and Caron, M.G. 2004. Hypertension and impaired glycine handling in mice lacking the orphan transporter XT2. *Mol. Cell. Biol.* 24: 4166-4173.
3. Höglund, P.J., Adzic, D., Scicluna, S.J., Lindblom, J. and Fredriksson, R. 2005. The repertoire of solute carriers of family 6: identification of new human and rodent genes. *Biochem. Biophys. Res. Commun.* 336: 175-189.
4. Romeo, E., Dave, M.H., Bacic, D., Ristic, Z., Camargo, S.M., Loffing, J., Wagner, C.A. and Verrey, F. 2006. Luminal kidney and intestine SLC6 amino acid transporters of BOAT-cluster and their tissue distribution in *Mus musculus*. *Am. J. Physiol. Renal Physiol.* 290: 376-383.
5. Eslami, B., Kinboshi, M., Inoue, S., Harada, K., Inoue, K. and Koizumi, A. 2006. A nonsense polymorphism (Y319X) of the solute carrier family 6 member 18 (SLC6A18) gene is not associated with hypertension and blood pressure in Japanese. *Tohoku J. Exp. Med.* 208: 25-31.
6. Kim, C.H., Waldman, I.D., Blakely, R.D. and Kim, K.S. 2008. Functional gene variation in the human norepinephrine transporter: association with attention deficit hyperactivity disorder. *Ann. N.Y. Acad. Sci.* 1129: 256-260.
7. Yoon, Y.H., Seol, S.Y., Heo, J., Chung, C.N., Park, I.H. and Leem, S.H. 2008. Analysis of VNTRs in the solute carrier family 6, member 18 (SLC6A18) and lack of association with hypertension. *DNA Cell Biol.* 27: 559-567

CHROMOSOMAL LOCATION

Genetic locus: SLC6A18 (human) mapping to 5p15.33.

PRODUCT

XTRP2 (h): 293T Lysate represents a lysate of human XTRP2 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

XTRP2 (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive XTRP2 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.