



DDB1 (h2): 293T Lysate: sc-116206

BACKGROUND

Damaged DNA binding protein (DDB) is a heterodimer composed of two subunits, p127 and p48, which are designated DDB1 and DDB2, respectively. The DDB heterodimer is involved in repairing DNA damaged by ultraviolet light. Specifically, DDB, also designated UV-damaged DNA binding protein (UV-DDB), xeroderma pigmentosum group E binding factor (XPE-BF) and hepatitis B virus X-associated protein 1 (XAP-1), binds to damaged cyclobutane pyrimidine dimers (CPDs). Mutations in the DDB2 gene are implicated as causes of xeroderma pigmentosum group E, an autosomal recessive disease in which patients are defective in nucleotide excision DNA repair. XPE is characterized by hypersensitivity of the skin to sunlight with a high frequency of skin cancer as well as neurologic abnormalities. The hepatitis B virus (HBV) X protein interacts with DDB1, which may mediate HBx transactivation.

REFERENCES

1. Dualan, R., Brody, T., Keeney, S., Nichols, A.F., Admon, A. and Linn, S. 1995. Chromosomal localization and cDNA cloning of the genes (DDB1 and DDB2) for the p127 and p48 subunits of a human damage-specific DNA binding protein. *Genomics* 29: 62-69.
2. Nichols, A.F., Ong, P. and Linn, S. 1996. Mutations specific to the xeroderma pigmentosum group E DDB-phenotype. *J. Biol. Chem.* 271: 24317-24320.
3. Stohr, H., Marquardt, A., Rivera, A., Kellner, U. and Weber, B.H. 1998. Refined mapping of the gene encoding the p127 kDa UV-damaged DNA-binding protein (DDB1) within 11q12-q13.1 and its exclusion in Best's vitelliform macular dystrophy. *Eur. J. Hum. Genet.* 6: 400-405.
4. Lin, G.Y., Paterson, R.G., Richardson, C.D. and Lamb, R.A. 1998. The V protein of the paramyxovirus SV5 interacts with damage-specific DNA-binding protein. *Virology* 249: 189-200.
5. Nichols, A.F., Itoh, T., Graham, J.A., Liu, W., Yamaizumi, M. and Linn, S. 2000. Human damage-specific DNA-binding protein p48. Characterization of XPE mutations and regulation following UV irradiation. *J. Biol. Chem.* 275: 21422-21428.
6. Zolezzi, F. and Linn, S. 2000. Studies of the murine DDB1 and DDB2 genes. *Gene* 245: 151-219.
7. Amundson, S.A., Do, K.T., Shahab, S., Bittner, M., Meltzer, P., Trent, J. and Fornace, A.J.J. 2000. Identification of potential mRNA biomarkers in peripheral blood lymphocytes for human exposure to ionizing radiation. *Radiat. Res.* 154: 342-346.
8. Wentz, M.J., Becker, S.A. and Slagle, B.L. 2000. Dissociation of DDB1-binding and transactivation properties of the hepatitis B virus X protein. *Virus Res.* 68: 87-92.
9. Wakasugi, M., Shimizu, M., Morioka, H., Linn, S., Nikaido, O. and Matsunaga, T. 2001. Damaged DNA-binding protein DDB stimulates the excision of cyclobutane pyrimidine dimers *in vitro* in concert with XPA and replication protein A. *J. Biol. Chem.* 276: 15434-15440.

CHROMOSOMAL LOCATION

Genetic locus: DDB1 (human) mapping to 11q12.2.

PRODUCT

DDB1 (h2): 293T Lysate represents a lysate of human DDB1 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

DDB1 (h2): 293T Lysate is suitable as a Western Blotting positive control for human reactive DDB1 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.