

ESX1 (h2): 293T Lysate: sc-116502

BACKGROUND

ESX1 was originally identified as a regulator of mouse embryogenesis. In mice, it is primarily expressed in placenta and testis where it functions in placenta/fetus development and spermatogenesis, respectively. In human cell lines, ESX1 has been elucidated as a paired-like homeoprotein that is proteolytically processed into N-terminal and C-terminal fragments. The N-terminal ESX1 fragment, which contains the homeodomain, localizes to the nucleus and represses mRNA transcription from the K-ras gene. A gain-of-function mutation of the K-ras gene is one of the most common genetic changes in human tumors. Therefore, ESX1 is implicated as a therapeutic target in the treatment of human cancers that have oncogenic K-ras mutations.

REFERENCES

1. Li, Y., Lemaire, P. and Behringer, R.R. 1997. ESX1, a novel X chromosome-linked homeobox gene expressed in mouse extraembryonic tissues and male germ cells. *Dev. Biol.* 188: 85-95.
2. Li, Y. and Behringer, R.R. 1998. ESX1 is an X-chromosome-imprinted regulator of placental development and fetal growth. *Nat. Genet.* 20: 309-311.
3. Yan, Y.T., Stein, S.M., Ding, J., Shen, M.M. and Abate-Shen, C. 2000. A novel PF/PN motif inhibits nuclear localization and DNA binding activity of the ESX1 homeoprotein. *Mol. Cell. Biol.* 20: 661-671.
4. Yanagihara, M., Ishikawa, S., Naito, M., Nakajima, J., Aburatani, H. and Hatakeyama, M. 2005. Paired-like homeoprotein ESXR1 acts as a sequence-specific transcriptional repressor of the human K-Ras gene. *Oncogene* 24: 5878-5887.
5. Yeh, Y.C., Yang, V.C., Huang, S.C. and Lo, N.W. 2005. Stage-dependent expression of extra-embryonic tissue-spermatogenesis-homeobox gene 1 (ESX1) protein, a candidate marker for X chromosome-bearing sperm. *Reprod. Fertil. Dev.* 17: 447-455.
6. Wang, X. and Zhang, J. 2007. Rapid evolution of primate ESX1, an X-linked placenta- and testis-expressed homeobox gene. *Hum. Mol. Genet.* 16: 2053-2060.
7. Nakajima, J., Ishikawa, S., Hamada, J., Yanagihara, M., Koike, T. and Hatakeyama, M. 2008. Anti-tumor activity of ESX1 on cancer cells harboring oncogenic K-Ras mutation. *Biochem. Biophys. Res. Commun.* 370: 189-194.

CHROMOSOMAL LOCATION

Genetic locus: ESX1 (human) mapping to Xq22.2.

PRODUCT

ESX1 (h2): 293T Lysate represents a lysate of human ESX1 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

ESX1 (h2): 293T Lysate is suitable as a Western Blotting positive control for human reactive ESX1 antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

ESX1 (B-9): sc-365740 is recommended as a positive control antibody for Western Blot analysis of enhanced human ESX1 expression in ESX1 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



ESX1 (B-9): sc-365740. Western blot analysis of ESX1 expression in non-transfected: sc-117752 (A) and human ESX1 transfected: sc-116502 (B) 293T whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.