

AIP2 (h2): 293T Lysate: sc-116898

BACKGROUND

Atrophin interacting proteins (AIPs) bind to atrophin-1 in the vicinity of the polyglutamine tract. The WW domain consists of 35-40 amino acids and is characterized by 4 well conserved aromatic residues, 2 of which are tryptophan. All five AIPs contain multiple WW domains and can be divided into two distinct classes. AIP1 and AIP3 (WWP3) are MAGUK-like multidomain proteins containing a guanylate kinase-like region, two WW domains and multiple PDZ domains. AIP2 (WWP2), AIP4 (itchy) and AIP5 (WWP1) are highly homologous, each having four WW domains and a HECT domain characteristic of ubiquitin ligases. These interactors are similar to isolated Huntingtin-interacting proteins, suggesting commonality of function between two families of proteins responsible for similar diseases.

REFERENCES

1. Bork, P. and Sudol, M. 1994. The WW domain: a signalling site in dystrophin? *Trends Biochem. Sci.* 19: 531-533.
2. André, B. and Springael, J.Y. 1994. WWP, a new amino acid motif present in single or multiple copies in various proteins including dystrophin and the SH3-binding Yes-associated protein YAP65. *Biochem. Biophys. Res. Commun.* 205: 1201-1205.
3. Hofmann, K. and Bucher, P. 1995. The Rsp5-domain is shared by proteins of diverse functions. *FEBS Lett.* 358: 153-157.
4. Pirozzi, G., McConnell, S.J., Uveges, A.J., Carter, J.M., Sparks, A.B., Kay, B.K. and Fowlkes, D.M. 1997. Identification of novel human WW domain-containing proteins by cloning of ligand targets. *J. Biol. Chem.* 272: 14611-14616.
5. Wood, J.D., Yuan, J., Margolis, R.L., Colomer, V., Duan, K., Kushi, J., Kaminsky, Z., Kleiderlein, J.J., Sharp, A.H. and Ross, C.A. 1998. Atrophin-1, the DRPLA gene product, interacts with two families of WW domain-containing proteins. *Mol. Cell. Neurosci.* 11: 149-160.
6. Perry, W.L., Hustad, C.M., Swing, D.A., O'Sullivan, T.N., Jenkins, N.A. and Copeland, N.G. 1998. The itchy locus encodes a novel ubiquitin protein ligase that is disrupted in a18H mice. *Nat. Genet.* 18:143-146.

CHROMOSOMAL LOCATION

Genetic locus: WWP2 (human) mapping to 16q22.1.

PRODUCT

AIP2 (h2): 293T Lysate represents a lysate of human AIP2 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

AIP2 (h2): 293T Lysate is suitable as a Western Blotting positive control for human reactive AIP2 antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

AIP2 (A-3): sc-398090 is recommended as a positive control antibody for Western Blot analysis of enhanced human AIP2 expression in AIP2 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



AIP2 (A-3): sc-398090. Western blot analysis of AIP2 expression in non-transfected: sc-117752 (A) and human AIP2 transfected: sc-116898 (B) 293T whole cell lysates.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.