

IP6K2 (h2): 293T Lysate: sc-117070

BACKGROUND

The members of the inositol hexakisphosphate kinase family, IP6K1 and IP6K2, have a high affinity and selectivity for inositol hexakisphosphate (InsP₆) as a substrate. IP6K1 and IP6K2 (also designated PiUS) convert InsP₆ to PP-InsP₅. However, neither kinase demonstrates any catalytic activity with other inositol pyrophosphates. The presence of InsP₆, which inhibits serine/threonine protein phosphatases, increases the influx of calcium across the plasma membrane and implies that it may mediate the regulation of Insulin exocytosis. IP6K1 was purified in rat brain extracts. By homology, IP6K1 and IP6K2 were characterized in mouse. IP6K1 displays ATP synthase activity by transferring a phosphate from PP-InsP₅ to ADP, which suggests a role for the IP6 kinases as high energy phosphate donors.

REFERENCES

1. Voglmaier, S.M., Bembenek, M.E., Kaplin, A.I., Dorman, G., Olszewski, J.D., Prestwich, G.D. and Snyder, S.H. 1996. Purified inositol hexakisphosphate kinase is an ATP synthase: diphosphoinositol pentakisphosphate as a high-energy phosphate donor. *Proc. Natl. Acad. Sci. USA* 93: 4305-4310.
2. Huang, C.F., Voglmaier, S.M., Bembenek, M.E., Saiardi, A. and Snyder, S.H. 1998. Identification and purification of diphosphoinositol pentakisphosphate kinase, which synthesizes the inositol pyrophosphate bis(diphospho) inositol tetrakisphosphate. *Biochemistry* 37: 14998-15004.
3. Saiardi, A., Erdjument-Bromage, H., Snowman, A.M., Tempst, P. and Snyder, S.H. 1999. Synthesis of diphosphoinositol pentakisphosphate by a newly identified family of higher inositol polyphosphate kinases. *Curr. Biol.* 9: 1323-1326.
4. Schell, M.J., Letcher, A.J., Brearley, C.A., Biber, J., Murer, H. and Irvine, R.F. 1999. PiUS (Pi uptake stimulator) is an inositol hexakisphosphate kinase. *FEBS Lett.* 461: 169-172.
5. Barker, C.J. and Berggren, P.O. 1999. Inositol hexakisphosphate and β -cell stimulus-secretion coupling. *Anticancer Res.* 19: 3737-3741.

CHROMOSOMAL LOCATION

Genetic locus: IP6K2 (human) mapping to 3p21.31.

PRODUCT

IP6K2 (h2): 293T Lysate represents a lysate of human IP6K2 transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

IP6K2 (h2): 293T Lysate is suitable as a Western Blotting positive control for human reactive IP6K2 antibodies. Recommended use: 10-20 μ l per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

IP6K2 (E-3): sc-390895 is recommended as a positive control antibody for Western Blot analysis of enhanced human IP6K2 expression in IP6K2 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



IP6K2 (E-3): sc-390895. Western blot analysis of IP6K2 expression in non-transfected: sc-117752 (A) and human IP6K2 transfected: sc-117070 (B) 293T whole cell lysates.



IP6K2 (G-9): sc-373770. Western blot analysis of IP6K2 expression in non-transfected: sc-117752 (A) and human IP6K2 transfected: sc-117070 (B) 293T whole cell lysates.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.