apoA-I (m): 293T Lysate: sc-118477



The Power to Question

BACKGROUND

Apolipoproteins are protein components of plasma lipoproteins. The human apoA-I gene encodes a single chain, 243 amino acid protein which promotes cholesterol efflux from tissues to the liver for excretion. apoA-I is the major protein component of high density lipoprotein (HDL) in the plasma. apoA-I can function as a cofactor for lecithin cholesterolacyltransferase (LCAT), which is responsible for the formation of most plasma cholesteryl esters. The human apoA-II gene encodes the second most abundant protein of HDL particles, where it influences plasma levels of free fatty acids. The human apoA-IV gene encodes a 396 amino acid preprotein, which after proteolytic processing is secreted from the intestine in association with chylomicron particles. ApoA-IV is a potent activator of LCAT *in vitro*. The human apoA-V gene encodes a 366 amino acid protein that is believed to be an important determinant of plasma triglyceride levels.

REFERENCES

- 1. Duriez, P. and Fruchart, J.C. 1999. High-density lipoprotein subclasses and apolipoprotein A-I. Clin. Chim. Acta 286: 97-114.
- Maezawa, I., et al. 2004. apoE isoforms and apoA-I protect from amyloid precursor protein carboxy-terminal fragment-associated cytotoxicity. J. Neurochem. 91: 1312-1321.
- 3. Maejima, T., et al. 2004. Effect of pitavastatin on apoA-I production in Hep G2 cell. Biochem. Biophys. Res. Commun. 324: 835-839.
- Maiorano, J.N., et al. 2004. Identification and structural ramifications of a hinge domain in apoA-I discoidal high-density lipoproteins of different size. Biochemistry 43: 11717-11726.
- Zhu, H.L., et al. 2004. Conformation and lipid binding of the N-terminal (1-44) domain of human apoA-l. Biochemistry 43: 13156-13164.
- 6. Cohen, J.C., et al. 2004. Multiple rare alleles contribute to low plasma levels of HDL cholesterol. Science 305: 869-872.
- 7. Fullerton, S.M., et al. 2004. The effects of scale: variation in the apoA-I/ C3/A4/A5 gene cluster. Hum. Genet. 115: 36-56.
- 8. Natarajan, P., et al. 2004. Identification of an apoA-I structural element that mediates cellular cholesterol efflux and stabilizes ATP binding cassette transporter A1. J. Biol. Chem. 279: 24044-24052.
- Kockx, M., et al. 2004. apoA-I-stimulated apoE secretion from human macrophages is independent of cholesterol efflux. J. Biol. Chem. 279: 25966-25977.

CHROMOSOMAL LOCATION

Genetic locus: APOA1 (human) mapping to 11q23.3.

PRODUCT

apoA-I (m): 293T Lysate represents a lysate of mouse apoA-I transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

apoA-I (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive apoA-I antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.


apoA-I (069-01): sc-58230 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse apoA-I expression in apoA-I transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-lgG κ BP-HRP: sc-516102 or m-lgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz MarkerTM Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA





apoA-I (069-01): sc-58230. Western blot analysis of apoA-I expression in non-transfected: sc-117752 (A) and mouse apoA-I transfected: sc-118477 (B) 293T whole cell I vsates.

apoA-I (2G4): sc-69755. Western blot analysis of apoA-I expression in non-transfected: sc-117752 (A) and mouse apoA-I transfected: sc-118477 (B) 293T

STORAGE

Store at -20 $^{\circ}$ C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3801 Fax 831.457.3801 Europe +00800 4573 8000 49 6221 4503 0 www.scbt.com