# Calpain 7 (m): 293T Lysate: sc-118965



The Power to Question

#### **BACKGROUND**

Calpains are calcium-activated thiol proteases. Calpain 7 (also known as PalBH) is a member of the non-EF-hand subfamily of Calpains and may be calcium-independent. Calpain 7 has 813 amino acid residues and is a divergent member of the Calpain family. It has only 26-35% shared identity to other members and most of this homology is in the protease domain. Calpain 7 seems to be related to PalB, an *Aspergillus nidulans* protease that is involved in alkaline ambient pH adaptation. A long N-terminal domain (N) and a PalB homologous domain (PBH) flank the Calpain protease domain of Calpain 7. Calpain 7 appears to have an ubiquitous tissue distribution, but is highly expressed in the brain. It localizes to the cytoplasm and the nucleus, but its activated form is found only in the nucleus. Calpain 7 is an atypical Calpain that lacks domain IV and cannot form a dimer with the 30 kDa regulatory subunit. Upregulation of Calpain 7 in striatal or cortical tissue of Huntington's disease knock-in mice suggests that this protein may be involved in the onset of the disease.

# **REFERENCES**

- 1. Takano, E., et al. 1991. Molecular diversity of erythrocyte calpastatin. Biomed. Biochim. Acta 50: 517-521.
- Takano, E., et al. 1993. Molecular diversity of calpastatin in human erythroid cells. Arch. Biochem. Biophys. 303: 349-354.
- 3. Johnson, G.V., et al. 1997. Calpains: intact and active? Bioessays 19: 1011-1018.
- Elce, J.S., et al. 1997. Autolysis, Ca<sup>2+</sup> requirement, and heterodimer stability in μ-Calpain. J. Biol. Chem. 272: 11268-11275.
- 5. Kawasaki, H., et al. 1997. Regulation of the Calpain-Calpastatin system by membranes (review). Mol. Membr. Biol. 13: 217-224.
- Dear, T.N., et al. 1999. Diverse mRNA expression patterns of the mouse Calpain genes Capn5, Capn6 and Capn11 during development. Mech. Dev. 89: 201-209.
- 7. Huang, Y., et al. 2001. The Calpain family and human disease. Trends Mol. Med. 7: 355-362.
- Gafni, J., et al. 2004. Inhibition of Calpain cleavage of Huntingtin reduces toxicity: accumulation of Calpain/caspase fragments in the nucleus. J. Biol. Chem. 279: 20211-20220.
- 9. Suzuki, K., et al. 2004. Structure, activation and biology of Calpain. Diabetes 53: S12-S18.

# **CHROMOSOMAL LOCATION**

Genetic locus: Capn7 (mouse) mapping to 14 B.

#### **PRODUCT**

Calpain 7 (m): 293T Lysate represents a lysate of mouse Calpain 7 transfected 293T cells and is provided as 100  $\mu$ g protein in 200  $\mu$ l SDS-PAGE buffer.

# **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

#### **APPLICATIONS**

Calpain 7 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive Calpain 7 antibodies. Recommended use: 10-20  $\mu$ l per lane

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

# **STORAGE**

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

# **PROTOCOLS**

See our web site at www.scbt.com for detailed protocols and support products.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3801 fax 831.457.3801 Europe +00800 4573 8000 49 6221 4503 0 www.scbt.com