SANTA CRUZ BIOTECHNOLOGY, INC.

Calponin 3 (m): 293T Lysate: sc-118971



BACKGROUND

Calponin regulates smooth muscle cell contraction and is a marker of smooth muscle cell differentiation. Calponin, an Actin- and Tropomyosin-binding protein, is characterized as an inhibitory factor of smooth-muscle actomyosin activity. Calponin is implicated in the regulation of smooth muscle contraction through its interaction with F-Actin and inhibition of the Actin-activated MgATPase activity of phosphorylated Myosin. Both properties are lost following phosphorylation (primarily at Serine 175) by protein kinase C or calmodulin-dependent protein kinase II. The three forms of Calponin, Calponin 1 (basic Calponin), Calponin 2 (neutral Calponin) and Calponin 3 (acidic Calponin) are found in smooth muscle tissue. Additionally, Calponin 2 is found in heart muscle tissue and Calponin 3 is found in the brain.

REFERENCES

- Tang, D.C., Kang, H.M., Jin, J.P., Fraser, E.D. and Walsh, M.P. 1996. Structure-function relations of smooth muscle Calponin. The critical role of serine 175. J. Biol. Chem. 271: 8605-8611.
- Masuda, H., Tanaka, K., Takagi, M., Ohgami, K., Sakamaki, T., Shibata, N. and Takahashi, K. 1996. Molecular cloning and characterization of human non-smooth muscle Calponin. J. Biochem. 120: 415-424.
- Doi, M., Kasuya, H., Weir, B., Cook, D.A. and Ogawa, A. 1997. Reduced expression of Calponin in canine basilar artery after subarachnoid haemorrhage. Acta Neurochir. 139: 77-81.
- Kaneko, T., Amano, M., Maeda, A., Goto, H., Takahashi, K., Ito, M. and Kaibuchi, K. 2000. Identification of Calponin as a novel substrate of Rho-kinase. Biochem. Biophys. Res. Commun. 273: 110-116.
- di Gioia, C.R., van de Greef, W.M., Sperti, G., Castoldi, G., Todaro, N., lerardi, C., Pieruzzi, F. and Stella, A. 2000. Angiotensin II increases calponin expression in cultured rat vascular smooth muscle cells. Biochem. Biophys. Res. Commun. 279: 965-969.
- Yoshimoto, R., Hori, M., Ozaki, H. and Karaki, H. 2000. Proteolysis of acidic Calponin by μ-Calpain. J. Biochem. 128: 1045-1049.

CHROMOSOMAL LOCATION

Genetic locus: Cnn3 (mouse) mapping to 3 G1.

PRODUCT

Calponin 3 (m): 293T Lysate represents a lysate of mouse Calponin 3 transfected 293T cells and is provided as 100 μg protein in 200 μl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

APPLICATIONS

Calponin 3 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive Calponin 3 antibodies. Recommended use: 10-20 μl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.