

β -casein (m5): 293T Lysate: sc-119013

BACKGROUND

Milk proteins are crucial for the development of all newborn mammals and caseins constitute the major proteins in mammalian milk. β - and κ -caseins are the only caseins present in human milk. The β -casein/ κ -casein ratio is higher in colostrum than in transitional and mature milk and is related to a better digestibility of colostrum casein micelles by the neonate during the first days of life. Human β -casein-encoding gene (Bca) contains a highly phosphorylated site, which is responsible for the calcium-binding capacity of β -casein. A common set of transcription factors are required for the expression of β -casein. Multiple binding sites for Stat5, C/EBP β (CCAAT/enhancer-binding protein) and several half-sites for glucocorticoid receptor (GR) are identified in the distal human enhancer of the β -casein gene. β -casein gene transcription is regulated primarily by a composite response element (CoRE), which integrates signaling from the lactogenic hormones PRL, Insulin and hydrocortisone in mammary epithelial cells. NF κ B functions as a negative regulator of β -casein gene expression during pregnancy by interfering with Stat5 tyrosine phosphorylation.

REFERENCES

- Greenberg, R., Groves, M.L. and Dower, H.J. 1984. Human β -casein. Amino acid sequence and identification of phosphorylation sites. J. Biol. Chem. 259: 5132-5138.
- Lonnerdal, B., Bergstrom, S., Andersson, Y., Hjalmarsson, K., Sundqvist, A.K. and Hernell, O. 1990. Cloning and sequencing of a cDNA encoding human milk β -casein. FEBS Lett. 269: 153-156.
- Menon, R.S., Chang, Y.F., Jeffers, K.F., Jones, C. and Ham, R.G. 1992. Regional localization of human β -casein gene (CSN2) to 4pter-q21. Genomics 13: 25-26.
- Hansson, L., Edlund, A., Johansson, T., Hernell, O., Stromqvist, M., Lindquist, S., Lonnerdal, B. and Bergstrom, S. 1994. Structure of the human β -casein encoding gene. Gene 139: 193-199.
- Winklehner-Jennwein, P., Geymayer, S., Lechner, J., Welte, T., Hansson, L., Geley, S. and Doppler, W. 1998. A distal enhancer region in the human β -casein gene mediates the response to prolactin and glucocorticoid hormones. Gene 217: 127-139.
- Cuilliere, M.L., Trégoat, V., Béné, M.C., Faure, G. and Montagne, P. 1999. Changes in the κ -casein and β -casein concentrations in human milk during lactation. J. Clin. Lab. Anal. 13: 213-218.
- Lykos, M.A., Fligger, J.M., Staley, M.D. and Baumrucker, C.R. 2000. Autocrine Insulin-like growth factor II inhibits β -casein mRNA expression in a mammary cell line. J. Dairy Sci. 83: 285-295.
- Wyszomierski, S.L. and Rosen, J.M. 2001. Cooperative effects of Stat5 (signal transducer and activator of transcription 5) and C/EBP β (CCAAT/enhancer-binding protein- β) on β -casein gene transcription are mediated by the glucocorticoid receptor. Mol. Endocrinol. 15: 228-240.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

CHROMOSOMAL LOCATION

Genetic locus: Csn2 (mouse) mapping to 5 E1.

PRODUCT

β -casein (m5): 293T Lysate represents a lysate of mouse β -casein transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

β -casein (m5): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive β -casein antibodies. Recommended use: 10-20 μ l per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

β -casein (H-4): sc-166530 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse β -casein expression in β -casein transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



β -casein (H-4): sc-166530. Western blot analysis of β -casein expression in non-transfected: sc-117752 (A) and mouse β -casein transfected: sc-119013 (B) 293T whole cell lysates.



β -casein (B-5): sc-393734. Western blot analysis of β -casein expression in non-transfected: sc-117752 (A) and mouse β -casein transfected: sc-119013 (B) 293T whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.