

Chk2 (m): 293T Lysate: sc-119229

BACKGROUND

Cell cycle events are regulated by the sequential activation and deactivation of cyclin dependent kinases (Cdks) and by proteolysis of cyclins. Chk1 and Chk2 are involved in these processes as regulators of Cdks. Chk1 and Chk2 both function as essential components in the G₂ DNA damage checkpoint by phosphorylating Cdc25C in response to DNA damage. Phosphorylation inhibits Cdc25C activity, thereby blocking mitosis. Cdc25A, Cdc25B and Cdc25C protein tyrosine phosphatases function as mitotic activators by dephosphorylating Cdc2 p34 on regulatory tyrosine residues. It has also been shown that Chk1 can phosphorylate Wee 1 *in vitro*, providing evidence that the hyperphosphorylated form of Wee 1, seen in cells delayed by Chk1 overexpression, is due to phosphorylation by Chk1.

REFERENCES

1. Gautier, J., et al. 1991. Cdc25 is a specific tyrosine phosphatase that directly activates p34 Cdc2. *Cell* 67: 197-211.
2. Barinaga, M. 1995. A new twist to the cell cycle. *Science* 269: 631-632.
3. O'Connell, M.J., et al. 1997. Chk1 is a Wee 1 kinase in the G₂ DNA damage checkpoint inhibiting Cdc2 by Y15 phosphorylation. *EMBO J.* 16: 545-554.
4. Sanchez, Y., et al. 1997. Conservation of the Chk1 checkpoint pathway in mammals: linkage of DNA damage to Cdk regulation through Cdc25. *Science* 277: 1497-1501.
5. Peng, C.Y., et al. 1997. Mitotic and G₂ checkpoint control: regulation of 14-3-3 protein binding by phosphorylation of Cdc25C on Serine 216. *Science* 277: 1501-1505.
6. Matsuoka, S., et al. 1998. Linkage of ATM to cell cycle regulation by the Chk2 protein kinase. *Science* 282: 1893-1897.
7. Zaugg, K., et al. 2007. Cross-talk between Chk1 and Chk2 in double-mutant thymocytes. *Proc. Natl. Acad. Sci. USA* 104: 3805-3810.
8. Abdelmohsen, K., et al. 2007. Phosphorylation of HuR by Chk2 regulates SIRT1 expression. *Mol. Cell* 25: 543-557.

CHROMOSOMAL LOCATION

Genetic locus: Chk2 (mouse) mapping to 5 F.

PRODUCT

Chk2 (m): 293T Lysate represents a lysate of mouse Chk2 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

Chk2 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive Chk2 antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

Chk2 (DCS-273): sc-56297 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse Chk2 expression in Chk2 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



Chk2 (DCS-273): sc-56297. Western blot analysis of Chk2 expression in non-transfected: sc-117752 (A) and mouse Chk2 transfected: sc-119229 (B) 293T whole cell lysates.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.