



Cofilin 2 (m): 293T Lysate: sc-119353

BACKGROUND

Cofilin 2, also known as CFL2 or Cofilin, muscle isoform, is a 166 amino acid protein that localizes to the cytoplasm and the cytoskeleton, as well as to the nuclear matrix, and contains one ADF-H domain. Existing as two alternatively spliced isoforms, one of which is known as CFL2 α and is expressed in heart and skeletal muscle, and the other of which is known as CFL2 β and is expressed ubiquitously, Cofilin 2 binds both F- and G-Actin and functions to control pH-dependent Actin polymerization and depolymerization. Cofilin 2 is subject to post-translational phosphorylation on Ser 24, an event which may prevent the nuclear localization of Cofilin 2. Defects in the gene encoding Cofilin 2 are the cause of nemaline myopathy type 7 (NEM7), a form of congenital myopathy that varies with age and is characterized by abnormal thread- or rod-like structures in muscle fibers.

REFERENCES

1. Gillett, G.T., Fox, M.F., Rowe, P.S., Casimir, C.M. and Povey, S. 1996. Mapping of human non-muscle type Cofilin (CFL1) to chromosome 11q13 and muscle-type Cofilin (CFL2) to chromosome 14. *Ann. Hum. Genet.* 60: 201-211.
2. Nebel, G., Meurer, S.C. and Samstag, Y. 1996. Dephosphorylation of serine 3 regulates nuclear translocation of Cofilin. *J. Biol. Chem.* 271: 26276-26280.
3. Bamburg, J.R., McGough, A. and Ono, S. 1999. Putting a new twist on Actin: ADF/Cofilins modulate Actin dynamics. *Trends Cell Biol.* 9: 364-370.
4. Thirion, C., Stucka, R., Mendel, B., Gruhler, A., Jaksch, M., Nowak, K.J., Binz, N., Laing, N.G. and Lochmüller, H. 2001. Characterization of human muscle type Cofilin (CFL2) in normal and regenerating muscle. *Eur. J. Biochem.* 268: 3473-3482.
5. Maciver, S.K. and Hussey, P.J. 2002. The ADF/Cofilin family: Actin-remodeling proteins. *Genome Biol.* 3: reviews3007.
6. Vartiainen, M.K., Mustonen, T., Mattila, P.K., Ojala, P.J., Thesleff, I., Partanen, J. and Lappalainen, P. 2002. The three mouse Actin-depolymerizing factor/Cofilins evolved to fulfill cell-type-specific requirements for Actin dynamics. *Mol. Biol. Cell* 13: 183-194.
7. Endo, M., Ohashi, K., Sasaki, Y., Goshima, Y., Niwa, R., Uemura, T. and Mizuno, K. 2003. Control of growth cone motility and morphology by LIM kinase and Slingshot via phosphorylation and dephosphorylation of Cofilin. *J. Neurosci.* 23: 2527-2537.
8. Agrawal, P.B., Greenleaf, R.S., Tomczak, K.K., Lehtokari, V.L., Wallgren-Pettersson, C., Wallefeld, W., Laing, N.G., Darras, B.T., Maciver, S.K., Dormitzer, P.R. and Beggs, A.H. 2007. Nemaline myopathy with minicores caused by mutation of the CFL2 gene encoding the skeletal muscle Actin-binding protein, Cofilin-2. *Am. J. Hum. Genet.* 80: 162-167.

CHROMOSOMAL LOCATION

Genetic locus: Cfl2 (mouse mapping to 12 C1).

PRODUCT

Cofilin 2 (m): 293T Lysate represents a lysate of mouse Cofilin 2 transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

Cofilin 2 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive Cofilin 2 antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.