

EDG-1 (m): 293T Lysate: sc-119919

BACKGROUND

The EDG (endothelial differentiation gene) family of G protein-coupled receptors consists of eight family members that bind lysophospholipid (LPL) mediators, including sphingosine-1-phosphate (SPP) and lysophosphatidic acid (LPA). EDG-1, EDG-3, EDG-5 (also designated H218 and AGR16) and EDG-8 bind SPP with high affinity. EDG-6 is a low affinity receptor for SPP. LPA preferentially binds to EDG-2, EDG-4 and EDG-7. The EDG receptors couple to multiple G proteins to signal through Ras, MAP kinase, Rho, Phospholipase C or other tyrosine kinases, which lead to cell survival, growth, migration and differentiation. EDG-1 signals through G_i proteins to activate Akt and is expressed in glioma cells. EDG-2 is expressed in brain, especially in white matter tract regions, while EDG-3 is expressed in cardiovascular tissue and in cerebellum. EDG-4 is highly expressed on leukocytes and brain, and EDG-5 has wide tissue distribution, including cardiovascular tissue and brain. EDG-6, which is expressed in lymphoid and hematopoietic tissues and in lung, signals through $G_{i/o}$ proteins, which activate growth related pathways.

REFERENCES

1. Goetzl, E.J. and An, S. 1999. A subfamily of G protein-coupled cellular receptors for lysophospholipids and lysosphingolipids. *Adv. Exp. Med. Biol.* 469: 259-264.
2. Van Brocklyn, J.R., Graler, M.H., Bernhardt, G., Hobson, J.P., Lipp, M. and Spiegel, S. 2000. Sphingosine-1-phosphate is a ligand for the G protein-coupled receptor EDG-6. *Blood* 95: 2624-2629.
3. Sato, K., Ui, M. and Okajima, F. 2000. Differential roles of EDG-1 and EDG-5, sphingosine 1-phosphate receptors, in the signaling pathways in C6 glioma cells. *Brain Res. Mol. Brain Res.* 85: 151-160.
4. Pyne, S. and Pyne, N.J. 2000. Sphingosine 1-phosphate signalling in mammalian cells. *Biochem. J.* 349: 385-402.
5. Zheng, Y., Kong, Y. and Goetzl, E.J. 2001. Lysophosphatidic acid receptor-selective effects on Jurkat T cell migration through a matrigel model basement membrane. *J. Immunol.* 166: 2317-2322.
6. Morales-Ruiz, M., Lee, M.J., Zoellner, S., Gratton, J.P., Scotland, R., Shiojima, I., Walsh, K., Hla, T. and Sessa, W.C. 2001. Sphingosine-1-phosphate activates Akt, nitric oxide production and chemotaxis through a G_i -protein/phosphoinositide 3-kinase pathway in endothelial cells. *J. Biol. Chem.* 276: 19672-19677.
7. Handford, E. J., Smith, D., Hewson, L., McAllister, G. and Beer, M.S. 2001. EDG-2 receptor distribution in adult rat brain. *Neuroreport* 12: 757-760.

CHROMOSOMAL LOCATION

Genetic locus: S1pr1 (mouse) mapping to 3 G1.

PRODUCT

EDG-1 (m): 293T Lysate represents a lysate of mouse EDG-1 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

EDG-1 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive EDG-1 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.