



Fibulin-2 (m): 293T Lysate: sc-120265

BACKGROUND

Fibulin-1 and Fibulin-2 associate with Fibronectin and other extracellular matrix proteins. In bone marrow, Fibulin-1 and Fibulin-2 bind to Fibronectin in the adherent layer. Fibulin-1 expression is stimulated by estrogen in ovarian cancer cell lines and has been suggested as both an agent of metastasis in ovarian cancer cells and as an indicator for predicting cancer risk or aggressiveness in ovarian carcinomas. The mobility of cancer cells may be inhibited with increasing exposure to Fibulin-1. Fibulin-2 binds to the lectin domains of extracellular matrix proteins aggrecan, versican and brevican. Fibulin-2 is abundantly expressed in heart, placenta and ovarian tissue, where it localizes to basement membranes and connective tissue compartments. In mice, differential Fibulin-2 gene expression correlates with the early phase of diabetic kidneys and glomerulosclerosis. The gene encoding human Fibulin-2 maps to chromosome 3p25.1.

REFERENCES

1. Argraves, W.S., et al. 1989. Fibulin, a novel protein that interacts with the Fibronectin receptor β -subunit cytoplasmic domain. *Cell* 58: 623-629.
2. Pan, T.C., et al. 1993. Structure and expression of Fibulin-2, a novel extracellular matrix protein with multiple EGF-like repeats and consensus motifs for calcium binding. *J. Cell Biol.* 123: 1269-1277.
3. Zhang, R.Z., et al. 1994. Fibulin-2 (Fbln-2): human cDNA sequence, mRNA expression, and mapping of the gene on human and mouse chromosomes. *Genomics* 22: 425-430.
4. Clinton, G.M., et al. 1996. Estrogens increase the expression of Fibulin-1, an extracellular matrix protein secreted by human ovarian cancer cells. *Proc. Natl. Acad. Sci. USA* 93: 316-320.
5. Roger, P., et al. 1998. Increased immunostaining of Fibulin-1, an estrogen-regulated protein in the stroma of human ovarian epithelial tumors. *Am. J. Pathol.* 153: 1579-1588.
6. Hayashido, Y., et al. 1998. Estradiol and Fibulin-1 inhibit motility of human ovarian- and breast-cancer cells induced by Fibronectin. *Int. J. Cancer* 75: 654-658.
7. Rochefort, H., et al. 1998. Estrogen receptor mediated inhibition of cancer cell invasion and motility: an overview. *Steroid Biochem. Mol. Biol.* 65: 163-168.
8. Gu, Y.C., et al. 2000. Association of extracellular matrix proteins Fibulin-1 and Fibulin-2 with fibronectin in bone marrow stroma. *Br. J. Haematol.* 109: 305-313.
9. Olin, A.I., et al. 2001. The proteoglycans aggrecan and versican form networks with Fibulin-2 through their lectin domain binding. *J. Biol. Chem.* 276: 1253-1261.

CHROMOSOMAL LOCATION

Genetic locus: Fbln2 (mouse) mapping to 6 D1.

PRODUCT

Fibulin-2 (m): 293T Lysate represents a lysate of mouse Fibulin-2 transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

Fibulin-2 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive Fibulin-2 antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.