

GCH-I (m2): 293T Lysate: sc-120451

BACKGROUND

GTP cyclohydrolase I (GCH-I), a homodecamer, catalyzes the conversion of GTP into dihydroneopterin triphosphate and folate. GCH-I is the first and rate limiting enzyme in tetrahydrobiopterin (BH4) biosynthesis. BH4 is the cofactor for tyrosine hydroxylase, a rate-limiting enzyme for dopamine synthesis and tryptophan hydroxylase, the rate-limiting enzyme for serotonin biosynthesis. Dopamine and serotonin are neurotransmitters involved in depression, which may be associated with a deficiency of BH4. Mutations in the gene encoding GCH-I cause malignant hyperphenylalaninemia, a genetic neurological disorder characterized by abnormally high levels of serum phenylalanine, and dopa-responsive dystonia (DRD), a group of movement disorders characterized by a progressive difficulty in walking which respond to L-dopa.

REFERENCES

1. Maita, N., et al. 2002. Crystal structure of the stimulatory complex of GTP cyclohydrolase I and its feedback regulatory protein GFRP. *Proc. Natl. Acad. Sci. USA* 99: 1212-1217.
2. Basset, G., et al. 2002. Folate synthesis in plants: the first step of the pterin branch is mediated by a unique bimodular GTP cyclohydrolase I. *Proc. Nat. Acad. Sci. USA* 99: 12489-12494.
3. Ishii, M., et al. 2005. Reduction of GTP cyclohydrolase I feedback regulating protein expression by hydrogen peroxide in vascular endothelial cells. *J. Pharmacol. Sci.* 97: 299-302.
4. Hagenah, J., et al. 2005. High mutation rate in dopa-responsive dystonia: detection with comprehensive GCH-I screening. *Neurology* 64: 908-911.
5. Hui xie, H., et al. 2005. Gene symbol: GTP cyclohydrolase I (GCH-I). Disease: dopa-responsive dystonia. *Hum. Genet.* 116: 235.
6. Nandi, M., et al. 2005. Pulmonary hypertension in a GTP-cyclohydrolase 1-deficient mouse. *Circulation* 111: 2086-2090.
7. Kealey, C., et al. 2005. Linkage and candidate gene analysis of 14q22-24 in bipolar disorder: support for GCH-I as a novel susceptibility gene. *Am. J. Med. Genet. B, Neuropsychiatr. Genet.* 136: 75-80.
8. Christensen, R., et al. 2005. Characterization of transgenic mice with the expression of phenylalanine hydroxylase and GTP cyclohydrolase I in the skin. *Exp. Dermatol.* 14: 535-542.
9. Pieper, G.M., et al. 2005. Nitric oxide formation in acutely rejecting cardiac allografts correlates with GTP cyclohydrolase I activity. *Biochem. J.* 391: 541-547.

CHROMOSOMAL LOCATION

Genetic locus: Gch1 (mouse) mapping to 14 C1.

PRODUCT

GCH-I (m2): 293T Lysate represents a lysate of mouse GCH-I transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

GCH-I (m2): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive GCH-I antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

GCH-I (G-8): sc-376483 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse GCH-I expression in GCH-I transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



GCH-I (G-8): sc-376483. Western blot analysis of GCH-I expression in non-transfected: sc-117752 (A) and mouse GCH-I transfected: sc-120451 (B) 293T whole cell lysates.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.