



group VI iPLA₂ (m): 293T Lysate: sc-120640

BACKGROUND

Phospholipases catalyze the release of fatty acids from phospholipids. One member of the phospholipase family, iPLA₂, is detected as a membrane-bound protein with multiple smaller isoforms, which result from alternative splicing. Two isoforms, ankyrin-iPLA₂-1 and -2, lack the catalytic domain and are thought to be involved in the negative regulation of iPLA₂ activity. The SH-iPLA₂ isoform is cytoplasmically localized. The human gene encoding iPLA₂ maps to chromosome 22q13.1. Another phospholipase, sPLA₂, belongs to a family of secretory phospholipases A₂, which represent an expanding family of related enzymes. sPLA₂ has both membrane bound and secreted forms that are encoded by a single gene. sPLA₂ is involved in the regulation of phospholipid metabolism in biomembranes and in eicosanoid biosynthesis.

REFERENCES

1. Scott, D.L., White, S.P., Browning, J.L., Rosa, J.J., Gelb, M.H. and Sigler, P.B. 1991. Structures of free and inhibited human secretory phospholipase A₂ from inflammatory exudate. *Science* 254: 1007-1010.
2. Lehninger, A., Nelson, A. and Cox, M. 1993. Principles of Biochemistry Second Edition. Worth Publishers.
3. Cupillard, L., Koumanov, K., Mattei, M.G., Lazdunski, M. and Lambeau, G. 1997. Cloning, chromosomal mapping, and expression of a novel human secretory phospholipase A₂. *J. Biol. Chem.* 272: 15745-15752.
4. Kitadokoro, K., Hagishita, S., Sato, T., Ohtani, M. and Miki, K. 1998. Crystal structure of human secretory phospholipase A₂-IIA complex with the potent indolizine inhibitor 120-1032. *J. Biochem.* 123: 619-623.
5. Ma, Z., Wang, X., Nowatzke, W., Ramanadham, S. and Turk, J. 1999. Human pancreatic islets express mRNA species encoding two distinct catalytically active isoforms of group VI phospholipase A₂ (iPLA₂) that arise from an exon-skipping mechanism of alternative splicing of the transcript from the iPLA₂ gene on chromosome 22q13.1. *J. Biol. Chem.* 274: 9607-9616.
6. Larsson-Forsell, P.K., Kennedy, B.P. and Claesson, H.E. 1999. The human calcium-independent phospholipase A₂ gene multiple enzymes with distinct properties from a single gene. *Eur. J. Biochem.* 262: 575-585.

CHROMOSOMAL LOCATION

Genetic locus: Pla2g6 (mouse) mapping to 15 E1.

PRODUCT

group VI iPLA₂ (m): 293T Lysate represents a lysate of mouse group VI iPLA₂ transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

group VI iPLA₂ (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive group VI iPLA₂ antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.